Antiepileptic drug prescription from studies prior to 2005

| Country (Ref) | Year | No of patients | Polytherapy% | VPA% | CBZ% | PHT% | PBT% | LTG% | TPM% | LEV% | OXC% | Others (%) |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Denmark (1) | 1998 | - | **26** | 28 | 9 | - | 1 | 24 | 2 | - | 24 | VGB(4), CLB(3), CZP (2), Primidone (1), TGB(0.5) |
| Netherlands (2) | 1996-2000 | - | **-** | 47 | 26 | 3 | - | 25 | - | - | - | - |
| Sweden (3) | 1996-2000 | 166 | **41** | 51 | 19 | - | - | 16 | - | - | - | CLB(10),CZP (7), STM (4), VGB(3), NZP(1), GBP(1), Felbamate(0.6), ETX(0.6) |
| Egypt (4) | 1999 | 229 | **22** | 38 | 48 | 31 | - | - | - | - | - | - |
| USA (5) | 1993-1997 | 613 | **-** | 18 | 39 | 11 | 10 | - | - | - | - | - |

References

1 Rochat P, Hallas J, Gaist D et al. Antiepileptic drug utilization: a Danish prescription database analysis. *Acta Neurol Scand.* 2001;104:6-11.

2 Knoester P, Belitser S, Deckers C et al. Diffusion of the new antiepileptic drug lamotrigine in Dutch clinical practice. *Eur J of Clin Pharmacol.* 2004;60:751-8.

3. Larsson K, Eeg-Olofsson O. A population based study of epilepsy in children from a Swedish county. *Europ J Paediatr Neurol.* 2006;10:107-13.

4. Bassili A, Omar T, Zaki A et al. Pattern of diagnostic and therapeutic care of childhood epilepsy in Alexandria, Egypt. *International Journal for Quality in Health Care.* 2002;14:277-84.

5. Berg AT, Levy SR, Testa FM et al. Treatment of newly diagnosed pediatric epilepsy: a community-based study. *Arch Pediatr Adolesc*. 1999;153:1267-71.