Laboratory reference intervals in the assessment of iron status in young children

Patricia C Parkin,^{1,2,3,4} Jemila Hamid,⁵ Cornelia M Borkhoff,^{1,2,3,4} Kawsari Abdullah,^{1,3} Eshetu G Atenafu,⁶ Catherine S Birken,^{1,2,3,4} Jonathon L Maguire,^{1,2,3,4,5,7} Azar Azad,^{8,9} Victoria Higgins,¹⁰ Khosrow Adeli^{9,10}

To cite: Parkin PC, Hamid J, Borkhoff CM, *et al*. Laboratory reference intervals in the assessment of iron status in young children. *BMJ Paediatrics Open* 2017;1:e000074. doi:10.1136/bmjpo-2017-000074

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjpo-2017-000074>).

Received 9 May 2017
Revised 30 July 2017
Accepted 7 August 2017

ABSTRACT

Objectives The primary objective was to establish reference intervals for laboratory tests used to assess iron status in young children using the Clinical and Laboratory Standards Institute guidelines. A secondary objective was to compare the lower limit of the reference interval with the currently recommended cut-off value for haemoglobin and serum ferritin in children 1–3 years of age.

Methods Blood samples were obtained from healthy children recruited during scheduled health supervision visits with their primary care physician. For our primary objective, outliers were removed; age partitions were selected and analysis of variance and pairwise comparisons were made between adjacent partitions; reference intervals and 90% CIs were calculated. For our secondary objective, we determined the proportion of children misclassified using the lower limit reference interval compared with the cut-off value.

Results Samples from 2305 male and 2029 female participants (10 days to 10.6 years) were used to calculate age and sex-specific reference intervals for laboratory tests of iron status. There were statistically significant differences between adjacent age partitions for most analytes. Approximately 10% of children 1–3 years of age were misclassified (underestimated) using the lower limit of the reference intervals rather than the currently recommended cut-off values for haemoglobin and serum ferritin.

Implications and relevance Clinical laboratories may consider adopting published paediatric reference intervals. Reference intervals may misclassify (underestimate) children with iron deficiency as compared with currently recommended cut-off values. Future research on decision limits derived from clinical studies of outcomes is a priority.

INTRODUCTION

Laboratory tests play an important role in clinical decision making.¹ Clinical laboratories are required to establish or verify ‘reference intervals’ (previously named ‘normal ranges’) for all tests offered by the laboratory.^{2,3} In conjunction with other health information, clinicians make diagnostic and treatment decisions based on their patients’ laboratory results in the context of the accompanying

What is already known on this topic?

- Iron deficiency is one of the most common single-nutrient deficiencies in both developing and developed countries, and peaks in prevalence in early childhood.
- For the laboratory assessment of iron status in young children, laboratory reference intervals are not well established and high-quality decision limits based on clinical studies are not available.

What this study hopes to add?

- This study presents age- and sex-specific reference intervals for haematological and biochemical tests commonly ordered by clinicians when assessing the iron status of children.
- Clinical laboratories may consider adopting the published paediatric reference intervals and age categories, rather than using adult reference intervals.
- Future research to develop high quality decision limits based on clinical studies of outcomes is a priority.

reference interval. However, clinicians are often unaware of the methods used to establish laboratory reference intervals. Reference intervals are derived from apparently healthy populations and provide two limits (lower and upper limits). The International Federation of Clinical Chemistry (IFCC) established the theory of reference intervals in 1987, and the Clinical and Laboratory Standards Institute (CLSI) provides guidelines for establishing and verifying reference intervals.^{4,5} These guidelines address issues of selection of the reference subjects, statistical methods including sample size requirements, approaches to grouping results according to categories such as age and sex (termed ‘partitions’) and approaches to the detection and removal of outliers.

CrossMark

For numbered affiliations see end of article.

Correspondence to

Patricia C Parkin; patricia.parkin@sickkids.ca

Reference intervals differ from 'decision limits'. Decision limits provide one (or more) cut-points intended to distinguish individuals with disease from those without disease.² According to a recently published quality hierarchy, decision limits based on clinical outcome studies are considered the highest quality; followed by reference intervals, and cut-off values based on clinicians' opinions or published professional recommendations.⁶ An example of a decision limit based on clinical outcome studies is HbA_{1c} for the diagnosis of diabetes.⁶

It has been recognised that establishing paediatric reference intervals poses specific challenges.^{3, 7} These challenges include the difficulties of obtaining samples from children, and the rapidly changing physiology from the newborn period to late adolescence. A recent systematic review of the current published literature examined the statistical methods used to construct paediatric reference intervals.⁸ Of the 22 publications identified, only 7 cited and followed the CLSI guidelines.

Iron deficiency is one of the most common single-nutrient deficiencies in both developing and developed countries, peaks in prevalence between 1 and 3 years of age and is associated with poor long-term neurodevelopment which may be irreversible.⁹ In a report on the diagnosis and prevention of iron deficiency in children under 3 years of age, the American Academy of Pediatrics (AAP) recommends universal screening for anaemia, with measurement of haemoglobin (Hb) concentration and consideration of additional tests to assess iron status, specifically serum ferritin (SF) and C-reactive protein (CRP). Although there are no established decision limits derived from clinical studies of outcomes, the AAP recommends a cut-off value for Hb of <110 g/L and for SF of <12 µg/L. The AAP describes other indicators of iron status including mean corpuscular volume, iron and transferrin. Few studies have examined the reference intervals for laboratory tests of iron status in young children, sample sizes have been small and age partitions wide.^{10–17}

The primary objective of this study was to establish reference intervals for haematological and biochemical tests commonly ordered by clinicians to assess the iron status in young children (10 years and younger) using the CLSI guidelines. A secondary objective was to compare the lower limit of the reference interval with the AAP currently recommended cut-off value for Hb and SF in children 1–3 years of age (age of peak prevalence for iron deficiency).

METHODS

Study population

Healthy children, from newborn to 10 years of age, were recruited during a scheduled health supervision visit with their primary care physician. Health supervision visits occur at 2 weeks, 2, 4, 6, 9, 12, 15, 18 months and then annually around the birthday of the child. Recruitment occurred between 2008 and 2016 at paediatric and family

medicine primary care practices participating in a practice-based research network called TARGet Kids! (www.targetkids.ca) in Toronto, Canada. The profile of this open longitudinal cohort has been previously described, and children with and without blood samples are similar with respect to demographics and health outcomes.¹⁸ Furthermore, the prevalence of iron deficiency is similar to other Canadian studies of this age group.¹⁹ Exclusion criteria were: children with associated health conditions affecting growth (eg, failure to thrive, cystic fibrosis), children with any acute or chronic conditions (other than asthma and high-functioning autism), children with severe developmental delay and families who are unable to communicate in English. Parents provided informed consent and completed a questionnaire regarding the child's age and sex.

Measures obtained by the TARGet Kids! research team include haematological and biochemical tests commonly ordered by clinicians to assess the nutritional status in young children. As a direct benefit of participating in the research, clinicians are provided with results of laboratory tests to be used for clinical purposes. For this study, we analysed data on haematological and biochemical markers of iron status recommended by the AAP; specifically Hb, mean corpuscular volume, three biochemical markers of iron status (SF, iron and transferrin) and CRP. As clinicians commonly request a complete blood count, other haematological tests, including white blood cell count and platelet count, were included in this analysis.

Sample collection and analysis

The research assistants, embedded in the primary care practices, are experienced phlebotomists and collected blood samples in lavender EDTA tubes and in serum separator tubes. Blood samples were refrigerated at the practice sites and transported to the laboratory at Mount Sinai Services (MSS) the same day. MSS provides services to researchers, is accredited and complies with regulatory standards (<http://www.mountsinaiservices.com/>). At the laboratory, blood samples were analysed fresh within 4–6 hours of collection. For haematology parameters, samples were analysed on the Sysmex XN-9000 Hematology Analyzer (Japan); for chemistry analytes, samples were analysed on the Roche platform (Switzerland) (see online supplementary table 1). The traceability chain used for SF was IS 80/602. The lower level of detection limit for CRP is 0.15 mg/L.

Statistical analysis

For our primary objective, data were analysed following CLSI C28-A3 guidelines and by choosing appropriate statistical methods based on sample size and distribution of the analytes.⁵ Data were first visually inspected using scatter plots, histograms and probability plots. Outliers were detected and removed using outlier detection techniques described by Horn *et al.*²⁰ Data were partitioned by sex (male/female). Age partitions were selected based on epidemiological knowledge of the changing

prevalence of iron deficiency and overall child growth and development. It is well established that 1–3 years is an age of peak prevalence for iron deficiency, thus we included a specific partition for this age group. Age partitions were as follows: <1 year (infant); >1 year to <3 years (toddler); >3 years to <6 years (preschool/early school age); >6 years to 10 years (school age). Analysis of variance (ANOVA) and pairwise comparisons between adjacent partitions were performed to determine if there was a difference on average laboratory measures across the different age partitions. In preparing the appropriate reference interval method, normality assumption, skewness as well as kurtosis were assessed for each of the laboratory measures per partition. Finally, the parametric or non-parametric method was used to estimate the reference interval as well as 90% CI around the reference intervals after excluding the outliers for each laboratory measure overall and within each partition. The parametric method was used if the normality assumption was met or the sample size in each partition was large; the non-parametric method was used if the normality assumption failed or the sample size was limited (see online supplementary table 2). All p values were two sided and for the statistical analyses, $p < 0.05$ indicated a significantly different result.

For our secondary objective, we further examined Hb and SF in the age partition of children 1–3 years of age, which is the age of peak prevalence for iron deficiency. Using the cut-off values recommended by the AAP (Hb <110 g/L; SF <12 µg/L), we determined the proportion of children misclassified using the lower limit reference interval compared with the cut-off values.

Statistical analysis was performed using SAS V.9.4 system for Windows (Copyright © 2002–2012 SAS Institute) and the R statistical software V.3.2.3 (2015. A language and environment for statistical computing, Vienna, Austria).

RESULTS

Samples from 2305 male and 2029 female participants (ages 10 days to 10.6 years) were used to calculate age and sex-specific reference intervals for haematological and biochemical tests of iron status. Age and sex are shown in table 1. Scatter plots for Hb and SF are shown in figures 1 and 2. Not all children had blood samples analysed for all analytes, resulting in different sample sizes for each analyte. Some children contributed a blood sample to more than one age group. Age and sex-specific paediatric

Table 1 Characteristics of study population—age and sex

Variable	Statistic	Summary
Sex	N (%)	
Male		2305 (53.2%)
Female		2029 (46.8%)
Age, months	Median (range)	36.6 (0.36–127)

Figure 1 Age-dependent scatter plots by sex of haemoglobin concentration (g/L). Red dots=male; blue dots=female.

reference intervals are shown in table 2. ANOVA across age partitions showed statistically significant differences for all analytes; adjacent partition pairwise comparisons of age partitions showed statistically significant differences for all analytes except for CRP (all age comparisons) and transferrin (3> to <6 years compared with ≥6 to 10 years).

For both Hb and SF in children 1–3 years of age, the lower limit of the reference interval was lower than the currently recommended AAP cut-off value. For Hb, the lower limit of the reference interval was 101 g/L (male) and 102 g/L (female), whereas the currently recommended AAP cut-off value is 110 g/L; the proportion misclassified (underestimated) was 10.78% for men and 9.76% for women. For SF, the lower limit of the reference interval was 6 µg/L (male) and 7 µg/L (female), whereas the currently recommended AAP cut-off value is 12 µg/L; the proportion misclassified (underestimated) was 10.19% for men and 10.97% for women.

DISCUSSION

We have established age and sex-specific reference intervals for several haematological and biochemical tests commonly ordered by clinicians when assessing the iron

Figure 2 Age-dependent scatter plots by sex of serum ferritin concentration (µg/L). Red dots=male; blue dots=female.

Table 2 Age and sex-specific reference intervals for analytes (after excluding outliers)

Haemoglobin, g/L						
Age group	Male			Female		
	N	Lower limit (90% CI)	Upper limit (90% CI)	N	Lower limit (90% CI)	Upper limit (90% CI)
10 days to <1 year	288	99 (97 to 100)	133 (132 to 134)	237	100 (99 to 102)	132 (130 to 133)
>1 to <3 years	815	101 (100 to 103)	133 (132 to 134)	707	102 (100 to 104)	134 (134 to 135)
>3 to <6 years	979	107 (106 to 108)	135 (134 to 136)	844	108 (107 to 109)	136 (135 to 136)
≥6 to 10 years	422	111 (108 to 112)	140 (139 to 141)	355	110 (109 to 111)	140 (139 to 141)
Mean corpuscular volume, fL						
10 days to <1 year	275	68 (65 to 70)	86 (84 to 88)	230	72 (71 to 73)	86 (85 to 86)
>1 to <3 years	814	67 (66 to 68)	83 (82 to 83)	687	69 (68 to 70)	83 (83 to 84)
>3 to <6 years	974	73 (72 to 73)	84 (84 to 85)	844	73 (72 to 74)	86 (85 to 86)
≥6 to 10 years	415	75 (75 to 76)	87 (86 to 87)	342	77 (75 to 77)	87 (86 to 88)
C-reactive protein, mg/L						
10 days to <1 year	296	0.09 (0.09 to 0.09)	6.10 (5.50 to 9.00)	243	0.09 (0.09 to 0.09)	4.70 (2.90 to 5.10)
>1 to <3 years	1001	0.04 (0.03 to 0.04)	3.091 (2.79 to 3.42)	896	0.09 (0.09 to 0.09)	7.30 (5.80 to 9.40)
>3 to <6 years	1195	0.043 (0.04 to 0.05)	3.00 (2.75 to 3.28)	1073	0.05 (0.05 to 0.06)	3.50 (3.18 to 3.84)
≥6 to 10 years	406	0.09 (0.09 to 0.09)	3.60 (2.70 to 4.10)	346	0.09 (0.09 to 0.09)	4.90 (4.00 to 8.90)
Serum ferritin, µg/L						
10 days to <1 year	266	9 (9 to 10)	115 (99 to 139)	233	10 (8 to 11)	194 (157 to 271)
>1 to <3 years	909	6 (5 to 6)	70 (65 to 79)	817	7 (6 to 8)	81 (75 to 89)
>3 to <6 years	1069	12 (11 to 12)	71 (68 to 74)	949	12 (11 to 12)	74 (71 to 81)
≥6 to 10 years	411	15 (13 to 15)	81 (71 to 87)	353	13 (10 to 15)	92 (83 to 109)
Iron, µmol/L						
10 days to <1 year	254	3 (2 to 4)	20 (18 to 21)	186	4 (3 to 5)	19 (17 to 20)
>1 to <3 years	786	4 (4 to 4)	21 (20 to 22)	705	4 (3 to 4)	23 (21 to 23)
>3 to <6 years	947	4 (4 to 5)	25 (23 to 26)	861	4 (4 to 4)	24 (23 to 25)
≥6 to 10 years	293	5 (4 to 5)	25 (24 to 27)	251	4 (3 to 5)	24 (23 to 26)
Transferrin, g/L						
10 days to <1 year	248	2.01 (1.95 to 2.08)	3.49 (3.40 to 3.54)	186	1.99 (1.92 to 2.06)	3.35 (3.28 to 3.43)
>1 to <3 years	765	2.32 (2.30 to 2.37)	3.80 (3.78 to 3.94)	669	2.30 (2.25 to 2.30)	3.70 (3.68 to 3.80)
>3 to <6 years	925	2.20 (2.20 to 2.27)	3.45 (3.40 to 3.50)	841	2.28 (2.20 to 2.30)	3.40 (3.39 to 3.46)
≥6 to 10 years	292	2.20 (2.17 to 2.28)	3.60 (3.40 to 3.64)	249	2.25 (2.15 to 2.35)	3.35 (3.30 to 3.42)
White blood cells, ×10⁹/L						
10 days to <1 year	293	6.51 (6.06 to 6.71)	17.0 (15.22 to 18.05)	236	5.81 (5.21 to 6.50)	16.02 (15.38 to 16.61)
>1 to <3 years	824	5.51 (5.14 to 5.89)	15.7 (15.1 to 16.1)	706	5.90 (5.42 to 6.00)	15.4 (14.8 to 15.9)
>3 to <6 years	988	4.59 (4.39 to 4.76)	12.9 (12.42 to 13.79)	859	4.70 (4.31 to 4.86)	12.7 (12.4 to 13.5)
≥6 to 10 years	420	4.47 (3.94 to 4.64)	12.0 (11.3 to 12.4)	351	4.47 (4.17 to 4.81)	12.7 (11.8 to 13.2)
Platelets, ×10⁹/L						
10 days to <1 year	281	262 (239 to 271)	598 (559 to 621)	228	237 (221 to 253)	573 (557 to 589)

Continued

Table 2 Continued

Age group	Haemoglobin, g/L					
	Male			Female		
	N	Lower limit (90% CI)	Upper limit (90% CI)	N	Lower limit (90% CI)	Upper limit (90% CI)
>1 to <3 years	816	200 (191 to 209)	513 (496 to 527)	706	213 (190 to 224)	530 (518 to 545)
>3 to <6 years	988	195 (184 to 204)	446 (441 to 465)	843	208 (201 to 214)	439 (432 to 443)
≥6 to 10 years	421	202 (190 to 210)	421 (414 to 435)	347	197 (184 to 202)	399 (389 to 442)

status of children. Age partitions showed statistically significant differences for most analytes, suggesting that laboratories should use these age categories. Furthermore, we found that approximately 10% of children 1–3 years of age would be misclassified as having healthy iron status (no disease) according to measures of Hb and SF, if the lower limit of the reference intervals was used rather

than the cut-off values currently recommended by the AAP.

In recognition of the critical gaps in paediatric reference intervals, the IFCC Task Force on Paediatric Laboratory Medicine launched an initiative to coordinate international efforts to fill this gap, representing groups from several countries.²¹ One group, the Canadian Laboratory Initiative on Pediatric Reference Intervals (CALIPER),

Table 3 CALIPER published reference intervals for several analytes relevant to the assessment of iron status in children

Analyte	Unit	Age range	Reference interval	Sample size, n	Reagent manufacturer	Study source	Reference
hsCRP	mg/L	0 to <14 days	0.3–6.1	139	Abbott	CALIPER	¹⁰
		15 days to <15 years	0.1–1.0	653			
Serum ferritin	µg/L	0 to <15 days	39.8–539.9	56	Beckman	CALIPER	¹⁵
		15 days to <6 months	15.3–374.6	49			
		6 months to <1 year	13.3–191.9	44			
		1 to <16 years	10.3–55.8	424			
		3–5 years	10.7–85.2	400	Ortho	CHMS	¹⁴
		Male 6–16 years	16.2–106.7	680			
		Female 6–24 years	9.6–81.9	790			
		4 to <15 days	99.6–717.0	127	Abbott	CALIPER	¹²
		15 days to <6 months	14.0–647.2	172			
		6 months to <1 year	8.4–181.9	140			
		1 to <5 years	5.3–99.9	126			
		5 to <14 years	13.7–78.8	497			
Haemoglobin	g/L	3–5 years	113.5–143.1	420	Beckman	CHMS	¹³
		6–8 years	114.7–143.0	670			
		9–10 years	118.4–146.9	540			
Iron	µmol/L	0 to <14 years	2.8–22.9	588	Abbott	CALIPER	¹⁰
MCV	fL	3–5 years	77.2–89.5	410	Beckman	CHMS	¹²
		6–11 years	77.8–91.1	1460			
Platelets	10 ⁹ /L	3–5 years	187.4–444.6	420	Beckman	CHMS	¹³
		6–9 years	186.7–400.4	910			
Transferrin	g/L	0 to <9 weeks	1.04–2.24	188	Abbott	CALIPER	¹⁰
		9 weeks to <1 year	1.07–3.24	104			
		1 to <19 years	2.20–3.37	878			

Reference values are applicable to both sexes, unless specified otherwise.

CALIPER, Canadian Laboratory Initiative on Pediatric Reference Intervals; CHMS, Canadian Health Measures Survey; hsCRP, high-sensitivity C-reactive protein; MCV, mean corpuscular volume.

has published several analyses that include reference intervals for several analytes relevant to the assessment of iron status in young children (see [table 3](#)); reference intervals are also accessible on their database website.^{10–15 22} However, data from CALIPER remain limited due to small sample sizes, especially for very young children, when iron deficiency peaks in prevalence. For example, Bailey *et al* presented reference intervals for SF based on 126 children 1–5 years of age.¹² Previous publications on the prevalence of iron deficiency in young children used data from the nationally representative US National Health and Nutrition Examination Survey (NHANES). These included a large number of laboratory tests for Hb, SF and other biochemical analytes.^{23–25} However, NHANES did not establish reference intervals according to the CLSI guidelines.

Given the challenges of developing reference intervals, it is difficult for each laboratory to develop their own reference intervals for children. As a result, clinicians are often provided with reference intervals for adults only which may lead to substantial misclassification, and possible overdiagnosis or underdiagnosis. Several prerequisites have been described in order for clinical laboratories to adopt common reference intervals.^{3 26} These include the following: the population served by the clinical laboratory is similar to that from which the reference interval was developed; the clinical laboratory uses methods that the manufacturers certify produce results that are traceable to the reference measurement system for a specific analyte; precision of the methods is within acceptable targets. It is also recommended that the clinical laboratory validate the reference interval in a small sample of their own population.^{3 27}

Reference intervals have inherent limitations in their application to clinical decision making. By their nature, reference intervals describe the characteristics of the referent population, which, while assumed to be healthy, may include asymptomatic individuals. Decision limits provide clinicians with the dichotomous information (disease/no disease) they seek for clinical decision making. In our study, we found wide reference intervals for SF, for example, which suggests that many children with asymptomatic iron deficiency were included in the referent population. Therefore, reference intervals may not adequately inform clinical decision making for the laboratory assessment of iron status in young children. Laboratories may also provide decision limits when available and cut-off values from published recommendations from professional organisations.⁶

Strengths of our study include a large sample of healthy children recruited from community settings during a health supervision visit with their primary care physician, and adherence to the CLSI guidelines. For the young age group at highest risk for iron deficiency (1–3 years), our sample size is substantially larger than previously published reference intervals.

Our study was limited to young children under 10 years of age. To fill the gaps in paediatric reference intervals,

other initiatives, such as CALIPER, have large sample sizes for children older than 10 years. Furthermore, we have not addressed cut-off values for children 3–10 years of age.

For our secondary objective, our study findings are limited by the low quality of evidence used to establish the currently recommended AAP cut-off value, especially for SF. The currently recommended cut-off value of 12 µg/L for SF is not based on a clinical outcome (as required for a high-quality decision limit), but rather based on the fifth percentile of the distribution of SF in a population presumed to be healthy.²³ In a recent analysis examining the relationship between Hb and SF, we identified a SF value of 23.7 µg/L corresponding to the level at which Hb concentration was maximised.²⁸ If the true decision limit for SF is indeed as high as 23.7 µg/L (rather than the currently recommended cut-off value of 12 µg/L), the proportion of children misclassified using the lower limit of the reference interval would be substantially higher.

CONCLUSION

We have established reference intervals for haematological and biochemical tests commonly ordered by clinicians to assess the iron status in young children under 10 years of age using the CLSI guidelines. As clinical laboratories are required to provide reference intervals for all tests offered, they may consider adopting published paediatric reference intervals. We have described the methodological approaches to establishing reference intervals and decision limits to better inform clinicians of their benefits and limitations as applied to clinical decision making. In the absence of decision limits based on clinical studies of outcomes (highest quality of evidence), we have described the currently recommended cut-off values for Hb and SF in the assessment of children 1–3 years of age and have identified the potential for misclassification when using reference intervals alone. Future research should focus on establishing decision limits for the laboratory assessment of iron status based on clinical outcomes.

Author affiliations

¹Division of Pediatric Medicine and the Pediatric Outcomes Research Team (PORT), Hospital for Sick Children, Toronto, Ontario, Canada

²Department of Pediatrics, Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada

³Child Health Evaluative Sciences, Hospital for Sick Children Research Institute, Toronto, Ontario, Canada

⁴Clinical Epidemiology and Health Care Research, Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, Ontario, Canada

⁵Li Ka Shing Knowledge Institute, St. Michael's Hospital, Toronto, Ontario, Canada

⁶Biostatistics Department, Princess Margaret Cancer Center, University Health Network, Toronto, Ontario, Canada

⁷Department of Pediatrics, St. Michael's Hospital, Toronto, Ontario, Canada

⁸Department of Pathology and Laboratory Medicine, Mount Sinai Hospital, Toronto, Ontario, Canada

⁹Department of Laboratory Medicine and Pathobiology, Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada

¹⁰CALIPER Program, Department of Pediatric Laboratory Medicine,, Hospital for Sick Children, Toronto, Ontario, Canada

Acknowledgements We thank all of the participating children and families for their time and involvement in the TARGet Kids! primary care practice-based research network and all practice site physicians, research staff, collaborating investigators, trainees, methodologists, biostatisticians, data management personnel, laboratory management personnel and advisory committee members (details may be found on our website: www.targetkids.ca).

Contributors PCP conceptualised and designed the study, designed the data collection instruments, interpreted the data, drafted the manuscript, critically revised and reviewed the manuscript for important intellectual content, and approved the final manuscript. JH conceptualised and designed the study, analysed and interpreted the data, critically revised and reviewed the manuscript for important intellectual content, and approved the final manuscript. CMB and KA conceptualised and designed the study, interpreted the data, critically reviewed the manuscript for important intellectual content and approved the final version of the manuscript. EGA performed the statistical analysis, interpreted the data, critically reviewed the manuscript for important intellectual content and approved the final version of the manuscript. JLM and CSB designed the data collection instruments, supervised the data collection, critically reviewed the manuscript for important intellectual content and approved the final version of the manuscript. AA, VH and KA interpreted the data, critically reviewed the manuscript for important intellectual content and approved the final version of the manuscript.

Funding Funding to support TARGet Kids! was provided by multiple sources including the Canadian Institutes for Health Research (CIHR), namely the Institute of Human Development, Child and Youth Health and the Institute of Nutrition, Metabolism and Diabetes, as well as the St. Michael's Hospital Foundation. The Pediatric Outcomes Research Team is supported by a grant from The Hospital for Sick Children Foundation. Funding agencies had no role in the design, collection, analyses or interpretation of the results of this study or in the preparation, review or approval of the manuscript.

Competing interests None declared.

Patient consent Obtained from the parents/guardian.

Ethics approval Hospital for Sick Children Research Ethics Board and the St. Michael's Hospital Research Ethics Board.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Piva E, Pelloso M, Penello L, *et al.* Laboratory critical values: automated notification supports effective clinical decision making. *Clin Biochem* 2014;47:1163–8.
- Cerioti F, Hinzmann R, Panteghini M. Reference intervals: the way forward. *Ann Clin Biochem* 2009;46(Pt 1):8–17.
- Jung B, Adeli K. Clinical laboratory reference intervals in pediatrics: the CALIPER initiative. *Clin Biochem* 2009;42:1589–95.
- Solberg HE. International Federation of Clinical Chemistry (IFCC), Scientific Committee, Clinical Section, Expert Panel on Theory of Reference Values, and International Committee for Standardization in Haematology (ICSH), Standing Committee on Reference Values. Approved Recommendation (1986) on the theory of reference values. Part 1. The concept of reference values. *J Clin Chem Clin Biochem* 1987;25:337–42.
- Clinical and Laboratory Standards Institute (CLSI). Defining, establishing, and verifying reference intervals in the clinical laboratory; approved guideline—third edition CLSI document C28-A.3; 2008.
- Sikaris K. Application of the Stockholm hierarchy to defining the quality of reference intervals and clinical decision limits. *Clin Biochem Rev* 2012;33:141–8.
- Cerioti F. Establishing pediatric reference intervals: a challenging task. *Clin Chem* 2012;58:808–10.
- Daly CH, Liu X, Grey VL, *et al.* A systematic review of statistical methods used in constructing pediatric reference intervals. *Clin Biochem* 2013;46:1220–7.
- Baker RD, Greer FR. Committee on Nutrition American Academy of Pediatrics. Diagnosis and prevention of iron deficiency and iron deficiency anemia in infants and young children (0–3 years of age). *Pediatrics* 2010;126:1040–50.
- Colantonio DA, Kyriakopoulou L, Chan MK, *et al.* Closing the gaps in pediatric laboratory reference intervals: a CALIPER database of 40 biochemical markers in a healthy and multiethnic population of children. *Clin Chem* 2012;58:854–68.
- Estey MP, Cohen AH, Colantonio DA, *et al.* CLSI-based transference of the CALIPER database of pediatric reference intervals from Abbott to Beckman, Ortho, Roche and Siemens Clinical Chemistry Assays: direct validation using reference samples from the CALIPER cohort. *Clin Biochem* 2013;46(13–14):1197–219.
- Bailey D, Colantonio D, Kyriakopoulou L, *et al.* Marked biological variance in endocrine and biochemical markers in childhood: establishment of pediatric reference intervals using healthy community children from the CALIPER cohort. *Clin Chem* 2013;59:1393–405.
- Adeli K, Raizman JE, Chen Y, *et al.* Complex biological profile of hematologic markers across pediatric, adult, and geriatric ages: establishment of robust pediatric and adult reference intervals on the basis of the Canadian Health Measures Survey. *Clin Chem* 2015;61:1075–86.
- Adeli K, Higgins V, Nieuwesteeg M, *et al.* Complex reference values for endocrine and special chemistry biomarkers across pediatric, adult, and geriatric ages: establishment of robust pediatric and adult reference intervals on the basis of the Canadian Health Measures Survey. *Clin Chem* 2015;61:1063–74.
- Karbasy K, Lin DC, Stoianov A, *et al.* Pediatric reference value distributions and covariate-stratified reference intervals for 29 endocrine and special chemistry biomarkers on the Beckman Coulter Immunoassay Systems: a CALIPER study of healthy community children. *Clin Chem Lab Med* 2016;54:643–57.
- Aldrimer M, Ridefelt P, Rödö P, *et al.* Population-based pediatric reference intervals for hematology, iron and transferrin. *Scand J Clin Lab Invest* 2013;73:253–61.
- Hilsted L, Rustad P, Aksglaede L, *et al.* Recommended Nordic paediatric reference intervals for 21 common biochemical properties. *Scand J Clin Lab Invest* 2013;73:1–9.
- Carsley S, Borkhoff CM, Maguire JL, *et al.* Cohort profile: The Applied Research Group for Kids (TARGet Kids!). *Int J Epidemiol* 2015;44:776–88.
- Hartfield D. Iron deficiency is a public health problem in Canadian infants and children. *Paediatr Child Health* 2010;15:347–50.
- Horn PS, Feng L, Li Y, *et al.* Effect of outliers and nonhealthy individuals on reference interval estimation. *Clin Chem* 2001;47:2137–45.
- Paediatric Reference Range Initiatives. <http://www.ifcc.org/task-force-paediatric-laboratory-medicine-web-pages/paediatric-reference-range-initiatives/> (accessed 20 Jan 2017).
- CALIPER Reference Interval Database. <http://www.sickkids.ca/caliperproject/intervals/> (accessed 20 Jan 2017).
- Looker AC, Dallman PR, Carroll MD, *et al.* Prevalence of iron deficiency in the United States. *JAMA* 1997;277:973–6.
- Cogswell ME, Looker AC, Pfeiffer CM, *et al.* Assessment of iron deficiency in US preschool children and nonpregnant females of childbearing age: National Health and Nutrition Examination Survey 2003–2006. *Am J Clin Nutr* 2009;89:1334–42.
- Gupta P, Perrine C, Mei Z, *et al.* Iron, Anemia, and Iron deficiency anemia among young children in the United States. *Nutrients* 2016;8:330.
- Cerioti F. Prerequisites for use of common reference intervals. *Clin Biochem Rev* 2007;28:115–21.
- Morrow DA, Cook NR. Determining decision limits for new biomarkers: clinical and statistical considerations. *Clin Chem* 2011;57:1–3.
- Abdullah K, Birken CS, Maguire JL, *et al.* Re-evaluation of serum ferritin cut-off values for the diagnosis of iron deficiency in children aged 12–36 months. *J Pediatr* 2017.