

Exploring pneumonia risk factors in Vietnamese infants: a survey of new mothers

Nguyen TK Phuong,^{1,2} Tran T Hoang,^{2,3} Kirsty Foster,^{4,5} Christine L Roberts,^{4,6} Ben J Marais^{1,4}

To cite: Phuong NTK, Hoang TT, Foster K, *et al.* Exploring pneumonia risk factors in Vietnamese infants: a survey of new mothers. *BMJ Paediatrics Open* 2017;1:e000155. doi:10.1136/bmjpo-2017-000155

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjpo-2017-000155>).

Received 11 June 2017
Revised 13 June 2017
Accepted 14 June 2017



CrossMark

¹Discipline of Paediatrics and Adolescent Medicine, The Children's Hospital at Westmead, The University of Sydney, Sydney, Australia

²Da Nang Hospital for Women and Children, Da Nang, Vietnam

³Da Nang University of Medical Technology and Pharmacy, Da Nang, Vietnam

⁴Sydney Medical School, The University of Sydney, Sydney, Australia

⁵Medical Education, Northern Clinical School & Kolling Institute, Sydney, Australia

⁶Clinical and Population Perinatal Health Research, Kolling Institute, Northern Sydney Local Health District, Sydney, Australia

Correspondence to

Dr Nguyen TK Phuong;
thng5150@uni.sydney.edu.au

ABSTRACT

Objective To understand the prevalence of pneumonia risk factors and perceived barriers to risk factor reduction among Vietnamese infants.

Methods We conducted a cross-sectional survey of mothers in the postnatal wards of Da Nang Hospital for Women and Children in central Vietnam from 10 February 2017 to 24 February 2017.

Results Of 286 mothers surveyed, 259 (91%) initiated breast feeding and 207 (72%) intended to continue exclusive breast feeding for 6 months. No mother smoked cigarettes, but 42% of fathers did. Mothers' decision not to smoke was motivated by concerns for their own health and that of their baby. Households rarely used wood or coal for cooking (6%). Mothers indicated near universal (99%) uptake of the National Expanded Program of Immunization vaccinations. Few (64; 22%) mothers knew about the pneumococcal conjugate vaccine; 56/64 (88%) indicated that they would purchase it for their infants. Family members rarely influenced mothers' decisions about breast feeding or vaccination, except in two instances where fathers were concerned about vaccine-related adverse effects.

Conclusion Modifiable pneumonia risk factors were uncommon among newborn babies in central Vietnam, apart from paternal cigarette smoke exposure. Successful local implementation of the WHO Essential Newborn Care package as well as high levels of maternal education and decision autonomy was observed.

BACKGROUND

Pneumonia is a common cause of death in young children worldwide.^{1 2} An estimated 900 000 children less than 5 years of age died because of pneumonia in 2015, with more than 90% of these deaths occurring in low-income and middle-income countries.³ In 2015, 14% of under-5 deaths in the Western Pacific region were attributed to pneumonia,⁴ with the majority (>75%) of pneumonia-related deaths occurring in six countries: Cambodia, China, Laos, Papua New Guinea, the Philippines and Vietnam.⁵ Despite recent progress, the pneumonia disease burden in Vietnam remains nearly 10 times higher than in

What is already known on this topic?

- Pneumonia is a common cause of morbidity and mortality in young children in Vietnam and worldwide.
- The WHO recognises lack of breast feeding, cigarette smoke or indoor air pollution exposure and vaccination as major modifiable risk factors for child pneumonia.
- Previous studies showed low rates of exclusive breast feeding, with high cigarette smoke and indoor air pollution exposure in Vietnam.

What this study hopes to add?

- We demonstrated a low prevalence of modifiable pneumonia risk factors, except for paternal cigarette smoke exposure, among newborn infants in central Vietnam.
- High breastfeeding initiation rates reflect recent successful implementation of the WHO Essential Newborn Care package at Da Nang Hospital for Women and Children.
- Mothers had high levels of education and displayed strong decision autonomy regarding what is best for their baby.

developed country settings like Australia and Europe.⁶ Efforts to reduce pneumonia-related morbidity and mortality in young children usually focus on vaccine-preventable causes and improved case management. In Vietnam, conjugated *Haemophilus influenzae* type B (Hib), measles, pertussis and BCG vaccination is provided free of charge by the National Expanded Program of Immunization (EPI). In general EPI uptake is excellent, although national Hib roll-out has only recently been achieved.^{4 7} The use of pneumococcal conjugate vaccine is advised in national guidance documents, but it is not provided free of charge in Vietnam and coverage is low.⁸

Apart from vaccination, primary prevention is rarely emphasised in settings that are dominated by income-generating healthcare

models, and often low on the priority list of developing countries with poor regulatory frameworks. The WHO recognises lack of breast feeding and cigarette smoke or indoor air pollution exposure as major modifiable risk factors for child pneumonia.^{9 10} In Vietnam, previous studies reported that less than 30% of mothers exclusively breast fed during the neonatal period¹¹ and less than 10% sustained exclusive breast feeding for up to 6 months of age.¹²⁻¹⁴ Although the health benefits of exclusive breast feeding, including significant reductions in infant pneumonia and death, are undisputed,¹⁵ many barriers compromise exclusive breast feeding during the first 6 months of life. Identified barriers include difficulties in establishing breast feeding, sociocultural factors and family expectations, as well as feasibility issues, inappropriate marketing of infant formula and poor health education.^{12 14 16} Studies from Vietnam listed 'working away from home', 'Caesarean section (C-section)' and 'maternal grandmother or husband preferences' as factors that reduce exclusive breast feeding.¹⁷⁻²⁰

Exposure to parental smoking increases the risk of severe pneumonia in children^{21 22} and is an independent risk factor of poor outcome.²³ In addition, indoor air pollution associated with the use of biomass fuel in homes is strongly associated with pneumonia-related mortality in children.^{24 25} Given the high burden of child pneumonia in Vietnam, there is a need to better understand the prevalence of pneumonia risk factors and perceived barriers to risk factor reduction, especially in central Vietnam where sociocultural practices are poorly documented.

METHODS

We conducted a prospective cross-sectional survey of mothers in the postnatal wards, within the first few days after delivery. The study was approved by the ethics review board at Da Nang Hospital for Women and Children (approved 26 August 2016) and performed over a 2-week period (10 February 2017 to 24 February 2017). All participating mothers provided written informed consent.

Study setting

The Da Nang Hospital for Women and Children is a regional hospital in central Vietnam with 1400 inpatient beds. It is the largest provider of obstetric services in Da Nang City with around 14000 deliveries each year. It also serves as the regional referral hospital for women and children. Pregnant women may come directly from home or be transferred from surrounding district and provincial hospitals to deliver their baby. The Da Nang Hospital for Women and Children recently implemented (fully implemented since November 2014) the WHO Essential Newborn Care (ENC) package,²⁶ which encourages early breastfeeding initiation.

Data collection and analysis

All mothers who were present in the postnatal wards when the researchers made their daily visits were recruited, if they provided informed consent to participate. Data were collected by the first author (NTKP) and a research assistant, neither of whom were involved in the clinical care of the mothers interviewed. Interviews were done in private without participation of staff members to limit possible social desirability bias. There were no exclusion criteria, and mothers were recruited irrespective of their date or mode of delivery (including vaginal deliveries and C-sections). However, recruitment only occurred in the general maternity wards, which effectively excluded mothers that required high-level care themselves or had infants that required neonatal intensive care. We recorded basic demographics (age, education level, address, type of delivery and reason for refusal) from mothers who declined participation or were otherwise unavailable for recruitment (not present in their rooms when the study team visited). Data were collected on a detailed questionnaire administered in Vietnamese by the investigator or trained assistant (see online supplementary questionnaire).

The questionnaire content was developed to focus on important modifiable risk factors for childhood pneumonia identified by the WHO. The questionnaire was translated into Vietnamese and back translated into English by a second independent translator to optimise the accuracy of the translation. A small group of Vietnamese mothers assisted questionnaire development to ensure that questions were correctly interpreted in the local cultural context. Further refinements were made after pilot testing was done on the maternity ward at Da Nang Hospital for Women and Children. Pilot testing occurred over a period of 3 days and involved more than 30 mothers. All questions were critically assessed and revised where necessary to improve efficiency, standardise data collection procedures and limit possible misinterpretation. Data quality checks were performed at the end of each day before entry into an Epidata database.²⁷ Descriptive analyses were performed using SPSS (V.24). Comparative analyses of factors that affected exclusive breast feeding will be presented separately.

RESULTS

In total, 573 mothers were admitted to the postnatal wards during the 2-week study period, of whom 302 were approached and 286 (53%) completed the postnatal questionnaire (figure 1). Only 16 (3%) mothers declined study participation, mostly because they were too tired to be interviewed. The age categories, occupation, mode of delivery and rural or urban origin were similar between study participants and those who were not enrolled (online supplementary file 2). All study participants were from the majority Kinh ethnic group.

Table 1 reflects the demographics of mothers who completed the questionnaire. Most (185; 65%) mothers




Figure 1 Study recruitment flow diagram.

were in the 25–34 age category; only one mother was less than 18 years and 16 (6%) were older than 40 years of age. Mothers were generally well educated, 64% attended college or university after high school completion and 55% were employed. Only one mother was HIV infected. Around half of mothers (55%) were primigravidas. **Table 2** describes the characteristics of the newborn infants and the healthcare that they received. The mean age of babies were 2.8 days (range: 1–5) with more boys than girls (55% vs 46%). More than half (162; 57%) were born by C-section, 55%) and 95% fell in the normal weight range; 5% were preterm.

Figure 2 provides an overview of common pneumonia risk factors reported during the interview. At the time of interview, 259 (91%) mothers had initiated breast feeding and 207 (72%) indicated that they intend to breast feed exclusively for 6 months. Not a single mother smoked cigarettes. Most mothers indicated consideration for their own health and that of their newborn baby as the main reasons for not smoking (**table 3**). By contrast, (120/286, 42%) of fathers smoked with a third (33/120, 28%) smoking more than 10 cigarettes per day; 23 (19%) smoked inside the house. We did not record data on other family members who may smoke in the house. Few families used wood (14, 5%) or charcoal (2, 1%) for cooking, and only 4 (1%) reported use of biomass fuel inside the house.

Table 4 summarises the mothers' knowledge of key pneumonia risk factors, with most mothers providing correct answers to basic questions on breast feeding and smoke exposure. As for vaccination, 284 (99%) mothers indicated that they intended adherence to the full EPI schedule. Of the non-EPI vaccines, 165 (58%) mothers were aware of the measles, mumps and rubella (MMR)

vaccine; 56% intended to provide this to their child. Only 64/286 (22%) mothers were aware of the pneumococcal conjugate vaccine and 56/286 (20%) intended to provide this to their infant. **Table 5** provides an overview of factors that influenced mothers' vaccination decisions. Mothers were mainly motivated by the fact that vaccines will protect their own babies, although there was awareness that "everyone should vaccinate their children" to protect the community. Only two mothers expressed fear of vaccine-related adverse effects and indicated that they will not vaccinate their children; their decisions were mostly influenced by concerns that their husbands expressed.

DISCUSSION

To our knowledge, this is the first study to explore the prevalence of pneumonia risk factors among newborn babies in central Vietnam. The rate of successful breast-feeding initiation and intent to continue exclusive breast feeding for 6 months among mothers was very high compared with national average estimates of 17% exclusive breast feeding for the first 6 months of life (in 2007–2014).⁴ The high rate of breastfeeding initiation is likely a reflection of recent adoption of the WHO Essential Newborn Care package at the Da Nang Hospital for Women and Children. The care package encourages healthcare professionals to assist mothers with breast feeding within the first hour after giving birth, together with regular skin-to-skin contact initiated at the time of delivery.²⁶ The hospital also provides mothers with knowledge on exclusive breast feeding through antenatal and postnatal consultation services.

Table 1 Demographics of mothers who completed the postpartum questionnaire

Demographics (n=286)	n (%)
Age (years)	
15–19	2 (1)
20–24	50 (18)
25–29	119 (41)
30–34	66 (23)
35–39	33 (11)
>40	16 (6)
Place of residence*	
Urban	156 (55)
Rural	130 (45)
Level of education†	
Primary	4 (1)
Secondary	32 (11)
High school	68 (24)
College	108 (38)
University	74 (26)
Occupation‡	
Employed	158 (55)
Self-employed	74 (26)
Unemployed	54 (19)
HIV infection	
Yes	3 (1)
No	283 (99)
Parity	
0 - First child	158 (55)
1 - Second child	88 (31)
2 - Third child	34 (12)
≥3 - Four or more children	6 (2)

*Place of residence: urban: administrative units (communes) that are part of recognised cities or large towns; rural: all other administrative units (communes).

†Level of education: primary: from years 1–5; secondary: from years 6–9; high school: from years 10–12; college/university: any qualification beyond high school.

‡Occupation: employed: have a paying job and will return to this job after maternity leave; self-employed: generate own income; unemployed: no income generation.

HIV, Human Immunodeficiency Virus.

The percentage of babies exposed to cigarette smoke at home was comparable with worldwide estimates (40% in 2011)²⁸ and slightly lower than national estimates for Vietnam (48% in 2012).¹⁰ The fact that only fathers were reported to smoke cigarettes concurs with high male smoking rates in Vietnam. In 2015, an estimated 36% of males in Vietnam smoked cigarettes compared with a global average of 25%.²⁹ It has been estimated that nearly a third (29%) of child pneumonia cases and 44 000 paediatric hospitalisation events in Vietnam can be

Table 2 Characteristics of babies born to mothers who completed the postpartum questionnaire

Characteristics (n=286)	n (%)*
Mean age (days)	2.8 (range 1–5)
Gender	
Boy	156 (55)
Girl	130 (45)
Gestational age†	
Preterm	15 (5)
Full-term	271 (95)
Post-term	0 (0)
Birth weight‡	
Low	11 (4)
Normal	261 (91)
High	14 (5)
Number of babies	
Singleton	283 (99)
Twins	3 (1)
Mode of delivery	
Caesarean section	162 (57)
Vaginal delivery	124 (43)
HBV vaccination	
Yes	263 (92)
Not yet	23 (8)
BCG vaccination§	
Not yet	286 (100)

*Unless otherwise specified.

†Preterm when the gestation age <37 weeks; full-term: 37–42 weeks; post-term: >42 weeks.

‡Low birth weight if weight <2500 g; normal weight: 2500–≤4000 g; overweight: >4000 g.

§BCG vaccine is normally not given to the baby at birth; infants receive BCG vaccine with their other routine vaccinations.

HBV, hepatitis B virus.

attributed to passive cigarette smoke exposure.³⁰ In utero exposure to cigarette smoke, or indoor air pollution, has also been associated with an increased risk of child pneumonia.^{25 31 32} However, studies showed that having a non-smoking mother significantly reduces the risk of hospitalisation from childhood pneumonia.³³ None of the mothers interviewed indicated that they smoke cigarettes, which is consistent with local clinical experience. The fact that mothers were motivated by concern for their babies health indicates that they could potentially also influence fathers to stop smoking. Although indoor air pollution is a major concern globally, very few families in our survey cooked with wood or charcoal and less than 2% burned biomass fuel inside the house. This figure is much lower than national solid fuel cooking rates of 47%, reported for Vietnam in 2013,⁴ which may be partly explained by the absence of minority ethnic groups from the study population.




Figure 2 Key pneumonia risk factors assessed during maternal postnatal interview. EPI: Expanded Program on Immunization; PCV: pneumococcal vaccine. Percentages were calculated based on the frequencies of each risk factors with 286 as nominator.

In the study setting, BCG vaccination was not provided in hospital, but in the community during the first month of life.³⁴ In countries with high rates of tuberculosis, WHO recommends that BCG vaccination should be provided at birth (before hospital discharge).^{34,35} Vietnam has a high tuberculosis incidence (137/100 000 population) indicative of ongoing community transmission, as well as high rates of drug resistant tuberculosis.³⁶ Infant BCG vaccination provides important protection against disseminated forms of disease, including drug-resistant tuberculosis, during the first 2–3 years of life.³⁷ There is room to consider whether BCG should best be administered before hospital discharge, but current arrangements are working well with good BCG uptake (95%, 2016)⁷ and low tuberculosis disease rates among young children.

Table 3 Reasons mothers identified for not smoking cigarettes

Reasons (n=286)	Most important n (%)	Second most important* n (%)
Smoking is bad for my own health	180 (63)	73 (34)
Smoking is bad for my baby's health	24 (8)	106 (49)
It is culturally unacceptable	80 (28)	37 (17)
Never considered smoking	2 (1)	NA

*This does not necessarily add up to N, since mothers had multiple options.

Additional pneumonia risk factors such as prematurity, low birth weight and HIV infection were uncommon in the study cohort; only one infant was born to an HIV-infected mother. The percentage of babies born by C-section was much higher than previous reports from Southern Vietnam (40%)²⁰ and other countries in the region (Cambodia 3%, Laos 4% and the Philippines 9%).⁴ Da Nang Hospital for Women and Children is a regional referral hospital with a more complex patient mix; however, C-section rates in Vietnam are generally higher than in neighbouring countries, which may

Table 4 Mothers' responses to questions on risk factors for child pneumonia

Question (n=286)	Answer		
	Yes n (%)	No n (%)	Unsure n (%)
Does breast feeding protect an infant against lung infections?*	246 (86)	21 (7)	19 (7)
Does cigarette smoke protect a child against lung infections?	10 (4)	274 (95)	2 (1)
Does smoke from cooking protect a child from lung infections?	18 (6)	261 (91)	7 (3)
Is it best to cook outside if cooking with charcoal or wood?	243 (85)	30 (11)	13 (4)

Correct answers in bold.

**Table 5** Factors that influenced mothers' vaccination choices for their babies

Factors		Most important n (%)	Second most important* n (%)
In mothers who were positive about immunisation (n=284)			
Reasons for vaccinating my child	Vaccines protect my baby	177 (62)	14 (18)
	It is the right thing to do†	65 (23)	51 (64)
	It is free of charge	37 (13)	14 (18)
	It is required to enter child care	1 (1)	1 (1)
	Other	4 (1)	NA
Main people who influenced my decision	I made the decision myself	173 (61)	13 (22)
	Health professional	74 (27)	21 (35)
	Husband	29 (10)	17 (28)
	Mother(s)	4 (1)	9 (15)
	Other	4 (1)	NA
In mothers who were negative about immunisation (n=2)			
Reasons for not vaccinating my child	I am scared of vaccine-related adverse effects	2 (100)	0 (0)
Main people who influenced my decision	Husband	2 (100)	NA

*This does not necessarily add up to N, since not all mothers indicated a second factor.

†Everyone should vaccinate their baby.

reflect differences in local clinical guidance and cultural expectations. It should be acknowledged that unnecessary C-section is a risk factor for neonatal respiratory distress and possible pneumonia.

Only two mothers indicated serious concerns regarding vaccine-related adverse effects. Most mothers were very keen to ensure that their children received the full EPI vaccination schedule. While many mothers planned to provide their children with MMR vaccination, in addition to measles vaccination provided by the EPI, few mothers had knowledge of newer pneumococcal conjugate vaccines. Of those who are new about pneumococcal conjugate vaccines, the majority planned to purchase this out of their own pocket. Vietnam has not yet indicated whether MMR or pneumococcal conjugate vaccines will be included in the state-sponsored EPI schedule. Previously documented barriers to vaccination uptake

include mothers from socioeconomic disadvantaged backgrounds, ethnic minorities, rural areas or those with limited education.³⁸ Of particular interest is the large degree of decision autonomy exercised by mothers, most of whom were primigravidas. Only a small minority indicated that family members or health professionals had a strong influence on their decisions to breast feed or vaccinate their children. Most mothers consulted with their husbands, mothers and healthcare professionals but indicated that they made the final decision themselves. The fact that mothers demonstrated a high degree of decision autonomy is encouraging and probably reflect their high education status and improved health literacy.

Our study was limited by the fact that we failed to interview a large number of mothers who delivered during study period. However, we purposefully guarded against selection bias and demonstrated comparable demographics among mothers who were interviewed and those not included in the study. Interviews were done within the first few days after delivery, and the hospital environment may have influenced mothers to provide the answers that they thought healthcare providers wanted. We tried to limit affirmation bias by framing all questions as objectively as possible and utilising study personnel who were not involved in the patient care. Our study did not include anyone from a minority ethnic group and therefore none of the study findings can be extrapolated to these populations, who may have increased exposure to indoor air pollution, lower health literacy and different barriers to vaccine uptake. It should also be emphasised that we only captured mothers' stated intent, without measuring their actual behaviour. However, we believe that the responses provided were truthful and provide useful insight into pneumonia risk factor exposure.

In conclusion, we documented a low prevalence of modifiable pneumonia risk factors, except for paternal cigarette smoke exposure, among newborn infants in central Vietnam, associated with high levels of maternal education and decision autonomy. High rates of breastfeeding initiation reflect recent successful implementation of the WHO Essential Newborn Care package at Da Nang Hospital for Women and Children.

Acknowledgements We would like to thank all the women who kindly agreed to participate, the staff at postnatal wards in Da Nang Hospital for Women and Children for supporting during the study period. Especially, we would like to thank mid-wife Le Thi Nhu Quynh and Tran Thi Xuan Oanh who helped with data collection.

Contributors All authors contributed to the design of the protocol and questionnaire. NTKP collected the data. NTKP and BJM conceptualised the manuscript and wrote the first draft. All authors gave constructive inputs and approved the final manuscript.

Funding No project funding received. Da Nang government provided an international PhD scholarship to the first author (NTKP).

Competing interests None declared.

Patient consent Obtained.

Ethics approval Da Nang Hospital for Women and Children Ethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Walker CLF, Rudan I, Liu L, *et al*. Theodoratou E, Bhutta ZqA, O'Brien KL, Campbell H, Black RE. Global burden of childhood pneumonia and diarrhoea. *Lancet* 2013;381:1405–16.
- Liu L, Johnson HL, Cousens S, *et al*. Child Health Epidemiology Reference Group of WHO, Unicef. Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000. *Lancet* 2012;379:2151–61.
- Liu L, Oza S, Hogan D, *et al*. Global, regional, and national causes of under-5 mortality in 2000–15: an updated systematic analysis with implications for the Sustainable Development Goals. *Lancet* 2016;388:3027–35.
- WHO. *World Health Statistics 2015*: World Health Organization, 2015.
- Naor B-Z, Carolyn M, John M, *et al*. Integrated Management of Childhood illness (IMCI) implementation in the Western Pacific Region: information package. *World Health Organ WPG*. 2013 http://www.wpro.who.int/child_adolescent_health/documents/imci_info_package/en/ (accessed 5 Feb 17).
- Nguyen TK, Tran TH, Roberts CL, *et al*. Child pneumonia - focus on the Western Pacific Region. *Paediatr Respir Rev* 2017;21:102–10.
- Viet Nam Ministry of Health. Report on vaccine uptake in EPI, Viet Nam from 2014–2016. 2017.
- World Health Organization. WHO vaccine - preventable diseases: monitoring system Global summary. 2016 http://apps.who.int/immunization_monitoring/globalsummary (accessed 25Mar 17).
- WHO. Pneumonia fact sheet no 331. www.who.int/mediacentre/factsheets/fs331/en/ (accessed 16 Jun).
- Nguyen TK, Tran TH, Roberts CL, *et al*. Risk factors for child pneumonia - focus on the Western Pacific Region. *Paediatr Respir Rev* 2017;21:95–101.
- WHO. *Workshop on Validation of Early Essential Newborn Care Progress*. Manila, Philippines: WHO Regional Office for the Western Pacific, 2015.
- Duong DV, Binns CW, Lee AH. Breast-feeding initiation and exclusive breast-feeding in rural Vietnam. *Public Health Nutr* 2004;7:795–9.
- Thu HN, Eriksson B, Khanh TT, *et al*. Breastfeeding practices in urban and rural Vietnam. *BMC Public Health* 2012;12:964.
- Duong DV, Lee AH, Binns CW. Determinants of breast-feeding within the first 6 months post-partum in rural Vietnam. *J Paediatr Child Health* 2005;41:338–43.
- Horta BL, Victora CG. *Short-Term effects of Breastfeeding: a Systematic Review on the Benefits of Breastfeeding on Diarrhea and Pneumonia Mortality*: WHO, 2013.
- Almroth S, Arts M, Quang ND, *et al*. Exclusive breastfeeding in Vietnam: an attainable goal. *Acta Paediatr* 2008;97:1066–9.
- Banapurmath CR, Selvamuthukumarasamy A. Initiation of breastfeeding in cesarean section mothers: antenatal advice versus postnatal assistance. *Indian Pediatr* 1995;32:902–5.
- Rowe-Murray HJ, Fisher JR. Baby friendly hospital practices: cesarean section is a persistent barrier to early initiation of breastfeeding. *Birth* 2002;29:124–31.
- Ramoo S, Trinh TA, Hirst JE, *et al*. Breastfeeding practices in a hospital-based study of Vietnamese women. *Breastfeed Med* 2014;9:479–85.
- Dearden KA, Quan leN, Do M, *et al*. Work outside the home is the primary barrier to exclusive breastfeeding in rural Viet Nam: insights from mothers who exclusively breastfed and worked. *Food Nutr Bull* 2002;23:99–106.
- Murtagh P, Cerqueiro C, Halac A, *et al*. Acute lower respiratory infection in argentinian children: a 40 month clinical and epidemiological study. *Pediatr Pulmonol* 1993;16:1–8.
- Karki S, Fitzpatrick AL, Shrestha S. Risk factors for Pneumonia in Children under 5 years in a Teaching Hospital in Nepal. *Kathmandu University Medical Journal* 2015;12:247–52.
- Jroundi I, Mahraoui C, Benmessaoud R, *et al*. Risk factors for a poor outcome among children admitted with clinically severe pneumonia to a university hospital in Rabat, Morocco. *Int J Infect Dis* 2014;28:164–70.
- Martin WJ, Glass RI, Balbus JM, *et al*. A Major environmental cause of death. *Science* 2011;334:180–1.
- Sonego M, Pellegrin MC, Becker G, *et al*. Risk factors for mortality from acute lower respiratory infections (ALRI) in children under five years of age in low and middle-income countries: a systematic review and meta-analysis of observational studies. *PLoS One* 2015;10:e0116380.
- WHO. Essential newborn care course. http://who.int/maternal_child_adolescent/documents/newborncare_course/en (accessed 4th Apr 17).
- EpiData Software. <http://www.epidata.dk/> accessed 27th May 17.
- Oberg M, Jaakkola MS, Woodward A, *et al*. Worldwide burden of disease from exposure to second-hand smoke: a retrospective analysis of data from 192 countries. *Lancet* 2011;377:139–46.
- GBD 2015 Tobacco Collaborators. Smoking prevalence and attributable disease burden in 195 countries and territories, 1990–2015: a systematic analysis from the global burden of disease study 2015. *Lancet* 2017;389:30819.
- Suzuki M, Thiem VD, Yanai H, *et al*. Association of environmental tobacco smoking exposure with an increased risk of hospital admissions for pneumonia in children under 5 years of age in Vietnam. *Thorax* 2009;64:484–9.
- Shibata T, Wilson JL, Watson LM, *et al*. Childhood acute respiratory infections and household environment in an eastern Indonesian urban setting. *Int J Environ Res Public Health* 2014;11:12190–203.
- Vanker A, Barnett W, Brittain K, *et al*. Antenatal and early life tobacco smoke exposure in an african birth cohort study. *Int J Tuberc Lung Dis* 2016;20:729–37.
- Barsam FJ, Borges GS, Severino AB, *et al*. Factors associated with community-acquired pneumonia in hospitalised children and adolescents aged 6 months to 13 years old. *Eur J Pediatr* 2013;172:493–9.
- WHO, Unicef. Recommended national immunization schedule for Viet Nam's children. 2015 http://wpro.who.int/vietnam/mediacentre/releases/2015/childhood_immunization_epi_schedule_march_2015_vietnam_v2.pdf (accessed 4th Apr 17).
- WHO. Weekly epidemiological record. *BCG vaccine* 2004;79:25–40.
- WHO. Global tuberculosis report 2016. 2016 http://www.who.int/tb/publications/global_report/en/ (accessed 5th Apr 17).
- Marais BJ, Seddon JA, Detjen AK, *et al*. WHO Child TB Subgroup. Interrupted BCG vaccination is a major threat to global child health. *Lancet Respir Med* 2016;4:251–3.
- Minh Anh D, Lee J-K, Van Minh H, *et al*. Timely immunization completion among children in Vietnam from 2000 to 2011: a multilevel analysis of individual and contextual factors. *Glob Health Action* 2016;9:29189.