



Questionnaire for the medical practitioner who requisitioned the EBV-tests:

| Was suspicion of mononucleosis the reason for taking the EBV-tests? | | | | □ YES □NO | | |
|---|---|--------|-------------------------|-----------|-----------------------------|-------|
| If yes | | | | | | |
| | Did the patient have fever? | | □ YES □ CE | | □DON'T KNOW | |
| | Did the patient have swollen lymph nodes of the neck? | | □ YES □ NO | | □DON'T KNOW | |
| | Did the patient have tonsillitis? | | \square YES \square | NO | □DON'T KNOW | |
| | Did the patient have pharyngitis? | | □ YES □ | NO | □DON'T KNOW | |
| Date for the first contact: | | Date | for estimated | l start | of disease: | |
| ` | was measured): patient's general practitioner? | | | _ ` | YES □NO | |
| If yes | Luncor Server Luncor | | | | | |
| ., | How long have you been the patient's GP? | | | | | years |
| How long h | ave you worked as a general practitioner/e | mergen | cy doctor? | _ | years | |
| you think tl | hat approx. 20% of all patients with monor his perticular patient will be among those 2 NO DON'T KNOW | | s are fatigued | d ½ y€ | ear after acute disease, do | |
| Other inform | mation: | | | | | |