



# Childhood immunisations in India during the COVID-19 pandemic

Anita Shet <sup>1</sup>, Baldeep Dhaliwal <sup>1</sup>, Preetika Banerjee,<sup>1</sup> Andrea DeLuca,<sup>2</sup> Kelly Carr,<sup>1</sup> Carl Britto,<sup>3</sup> Rajeev Seth,<sup>4</sup> Bakul Parekh,<sup>5</sup> Gangasamudra V Basavaraj,<sup>5</sup> Digant Shastri,<sup>5</sup> Piyush Gupta<sup>5</sup>

**To cite:** Shet A, Dhaliwal B, Banerjee P, *et al.* Childhood immunisations in India during the COVID-19 pandemic. *BMJ Paediatrics Open* 2021;**5**:e001061. doi:10.1136/bmjpo-2021-001061

Received 13 February 2021  
Revised 25 March 2021  
Accepted 5 April 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>International Vaccine Access Center, Department of International Health, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland, USA

<sup>2</sup>Amputee Coalition of America, Knoxville, Tennessee, USA

<sup>3</sup>Unit of Infectious Diseases, St John's Research Institute, Bangalore, Karnataka, India

<sup>4</sup>Bal Umang Drishya Sanstha (BUDS), New Delhi, India

<sup>5</sup>Indian Academy of Pediatrics, Navi Mumbai, Maharashtra, India

## Correspondence to

Dr Anita Shet; ashet1@ju.edu

The COVID-19 pandemic has led to major disruptions in the delivery of essential health services including routine immunisation services in many countries, setting the stage for potentially serious population health effects. The WHO reported major disruptions to vaccination services in countries around the world, and estimated that approximately 80 million children under the age of 1 were living in countries where routine immunisation services were disrupted and could potentially be at risk of developing a vaccine-preventable illness.<sup>1</sup> Evidence from previous epidemics has demonstrated that even temporary interruptions of routine immunisation services can lead to secondary public health crises, such as outbreaks of vaccine-preventable diseases, amplifying morbidity and mortality.<sup>2</sup> This commentary explores the possible effects of the COVID-19 pandemic on routine immunisations in India.

Results from a survey of Indian paediatricians<sup>3</sup> amplify a growing chorus around the globe calling for a focus on vaccine-preventable illnesses, even as COVID-19 cases grow worldwide. This comes at a pivotal time in India's ongoing pursuit to improve immunisation coverage. The national immunisation programme run by the Government of India is one of the largest in the world, with an annual reach of over 26 million children and 29 million pregnant women.<sup>4</sup> Mission Indradhanush was launched in 2014 to extend this reach and achieve full immunisation for 90% of children, and the programme was further intensified in 2019.<sup>4 5</sup> While remarkable progress has been made, there is evidence of existing inequalities in coverage.<sup>6</sup>

Early in the pandemic, soon after the lockdown was announced, there were major disruptions in health services, especially in women and children's services. Movement restrictions were likely to have disrupted strategies used by Mission Indradhanush,

including community mobilisation, door-to-door campaigns and monitoring events. The National Health Mission's health management and information system reported a substantial decrease in routine immunisation services relative to the previous year, indicating that in March 2020 at least 100 000 and 200 000 children missed their BCG and pentavalent (diphtheria, tetanus, pertussis, hepatitis B and *Haemophilus influenzae* type b) vaccines, respectively.<sup>7 8</sup> Researchers modelled different scenarios and used the Lives Saved Tool (LiST, a mathematical modelling tool to estimate the impact of programme coverage of maternal, newborn and child health on mortality at a country level<sup>9</sup>) to demonstrate that widespread disruption to health systems could lead to substantial increases in maternal and child deaths.<sup>10</sup> In India they estimated that an additional 49 000 child deaths and 2300 maternal deaths in a month could be attributable to severely disrupted services.<sup>10</sup> Applying the current population demographic data in India, estimates suggested that eventually over 27 million children will miss out on diphtheria tetanus pertussis vaccines and other health services, resulting in a 40% increase in child mortality over the next year.<sup>11</sup> If vaccination services are not restored and barriers to access are not addressed, disparities will become more pronounced and the number of zero-dose children will likely increase. India currently accounts for 2.1 million of the 20 million unvaccinated and undervaccinated children globally (11%),<sup>12</sup> and the national lockdown has shown its potential to further exacerbate this problem. Without targeted campaigns and effort, there is a legitimate risk for a reversal of gains made through national programmes.

Experiences from past outbreaks provide lessons on the indirect impacts which can be even more harmful to health. Analyses from the West African Ebola outbreak in 2014–2015



suggest that the increased number of deaths caused by other infections such as measles, HIV and tuberculosis attributable to health system failures exceeded deaths from Ebola.<sup>2 13</sup> A sustained period of disrupted immunisation can result in the accumulation of susceptible individuals, which in turn can lead to disease outbreaks.<sup>14</sup> Given the disruptions and the realisation of the dire consequences, the Government of India declared immunisation an essential health service in April 2020 and issued guidelines for states to resume routine immunisation services.<sup>15</sup> In June 2020 India began a phased reopening of the economy, and the resumption of immunisation activities was appropriately structured based on local COVID-19 infection rates and restrictions. These activities were based on the WHO guidance urging nations to continue providing essential services along with COVID-19 mitigation and treatment measures in order to maintain public trust and minimise morbidity and mortality.<sup>16</sup>

Coordinated campaigns across India targeting children who missed critical routine vaccinations during the national lockdown, as well as targeting low-coverage areas, could prevent additional public health disasters. Prioritising measles vaccine catch-up would be most prudent given the outbreak potential with even marginal reduction in herd immunity.<sup>17</sup> Planning catch-up campaigns now is essential so providers can minimise the time children are at risk of vaccine-preventable diseases. Vaccination catch-up sessions could institute innovative strategies such as implementing appointment-only visits or designated walk-in clinics for healthy children, minimising overcrowding, separating immunisation visits from sick children visits, prioritising robust communication efforts which address caregivers' fears of contracting COVID-19, and sending reminders to caregivers of the importance of routine vaccinations.<sup>18</sup> In addition, healthcare strategies such as the Integrated Management of Newborn and Childhood Illnesses can strengthen their focus on immunisation. Empowering community health workers to trace children who missed vaccination appointments can help restore baseline vaccination levels. Catch-up vaccinations can also be given to children in contact healthcare facilities for acute or chronic illnesses.<sup>19</sup> There is growing evidence that the risk to benefit ratio is decidedly in favour of continuing vaccination services even when considering the consequences of doing so during the pandemic.<sup>20</sup> Gaining provider insights on effective strategies is essential to establishing context-specific mechanisms to prioritise catch-up for missed vaccines. In neighbouring country Pakistan, an analysis of predictors associated with immunisation during their lockdown showed that factors such as higher maternal education, facility-based births and early enrolment into the immunisation programme were associated with higher immunisation uptake.<sup>21</sup> Interventions targeted at sustaining these predictors could be effective means of engaging with caregivers to ensure catch-up for missed vaccines in India as well. A pulse survey from the WHO indicated

partial disruption of essential health services beyond immunisations in many regions of the world, particularly in lower-income countries,<sup>22</sup> which prompted a strong call to arms for health systems and governments to 'build back better' to incorporate health system resilience and maintain the provision of essential health services during and after the COVID-19 pandemic.<sup>23</sup>

The Government of India has recently worked on several strategies for health systems strengthening, including incorporating a coordinated programme for public health surveillance, which will help monitor outbreaks of vaccine-preventable diseases among other diseases.<sup>24</sup> In the context of the pandemic, the Government of India has approved COVID-19 vaccines and the nation has embarked on one of the largest and most ambitious immunisation campaigns in the world. Although children will not receive the COVID-19 vaccine at this time, their caregivers and healthcare providers who will receive the vaccine should be provided with targeted messages and reminders for childhood routine immunisations. In addition, liaising routine immunisation campaigns with the COVID-19 vaccine roll-out, particularly in hard-to-reach areas, would be beneficial, given India's vast human resources and immunisation experience. Concerted efforts are needed from governing and academic groups to ensure that routine immunisation and catch-up programmes are implemented to sustain gains in vaccination coverage and provide a robust blueprint for the national roll-out of the COVID-19 vaccine.

**Contributors** AS prepared the first draft of the manuscript. BD, PB, ADL and DS provided critical review and input. All authors reviewed the manuscript and approved the final version for submission.

**Funding** This work was supported by the Johns Hopkins Maternal and Child Health Center, India, and the Indian Academy of Pediatrics.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Anita Shet <http://orcid.org/0000-0002-7204-8164>

Baldeep Dhaliwal <http://orcid.org/0000-0001-5021-3432>

#### REFERENCES

- World Health organization, news: at least 80 million children under one at risk of diseases such as diphtheria, measles and polio as COVID-19 disrupts routine vaccination efforts, warn Gavi, who and UNICEF. Available: <https://www.who.int/news/item/22-05-2020-at-least-80-million-children-under-one-at-risk-of-diseases-such-as-diphtheria-measles-and-polio-as-covid-19-disrupts-routine-vaccination-efforts-warn-gavi-who-and-unicef> [Accessed 22 May 2020].
- Takahashi S, Metcalf CJE, Ferrari MJ, *et al*. Reduced vaccination and the risk of measles and other childhood infections post-Ebola. *Science* 2015;347:1240-2.

- 3 Shet A, Dhaliwal B, Banerjee P. COVID-19-related disruptions to routine vaccination services in India: perspectives from pediatricians. *medRxiv* 2021.
- 4 Gurnani V, Halder P, Aggarwal MK, *et al*. Improving vaccination coverage in India: lessons from intensified mission Inradhanush, a cross-sectoral systems strengthening strategy. *BMJ* 2018;363:k4782.
- 5 Intensified mission Inradhanush 2.0: coverage report. 25 September 2020. Available: <https://imi2.nhp.gov.in/report/coverage> [Accessed 20 Nov 2020].
- 6 Srivastava S, Fledderjohann J, Upadhyay AK. Explaining socioeconomic inequalities in immunisation coverage in India: new insights from the fourth national family health survey (2015-16). *BMC Pediatr* 2020;20:295.
- 7 National health mission health management information system, Ministry of health and family welfare, government of India. Available: <https://nrhm-mis.nic.in/SitePages/Home.aspx> [Accessed on 15 Nov 2020].
- 8 Rukmini S. How covid-19 response disrupted health services in rural India. *mint*, 2020. Available: <https://www.livemint.com/news/india/how-covid-19-response-disrupted-health-services-in-rural-india-11587713155817.html> [Accessed on 15 Nov 2020].
- 9 The lives saved tool. Available: <https://www.livessavedtool.org/>
- 10 Robertson T, Carter ED, Chou VB, *et al*. Early estimates of the indirect effects of the COVID-19 pandemic on maternal and child mortality in low-income and middle-income countries: a modelling study. *Lancet Glob Health* 2020;8:e901–8.
- 11 Global Financing Facility. Preserve essential health services during the COVID-19 pandemic: India, 2020. Available: [https://idronline.org/user-content/uploads/2020/06/India-Covid-Brief\\_GFF.pdf](https://idronline.org/user-content/uploads/2020/06/India-Covid-Brief_GFF.pdf) [Accessed 18 Nov 2020].
- 12 Progress and challenges with achieving universal immunisation coverage. 2019 WHO/UNICEF estimates of national immunisation coverage. Available: [https://www.who.int/immunisation/monitoring\\_surveillance/who-immuniz.pdf](https://www.who.int/immunisation/monitoring_surveillance/who-immuniz.pdf) [Accessed 15 Jul 2020].
- 13 Elston JWT, Cartwright C, Ndumbi P, *et al*. The health impact of the 2014-15 Ebola outbreak. *Public Health* 2017;143:60–70.
- 14 Truelove SA, Graham M, Moss WJ, *et al*. Characterizing the impact of spatial clustering of susceptibility for measles elimination. *Vaccine* 2019;37:732–41.
- 15 Immunisation services during and post COVID-19 outbreak. Ministry of health and family welfare, government of India. Available: <https://www.mohfw.gov.in/pdf/3/ImmunisationServicesduringCOVIDOutbreakSummary150520202.pdf> [Accessed 15 Apr 2020].
- 16 World Health Organization. Maintaining essential health services: operational guidance for the COVID-19 context interim guidance. Available: <https://www.who.int/publications/i/item/covid-19-operational-guidance-for-maintaining-essential-health-services-during-an-outbreak> [Accessed 18 Nov 2020].
- 17 Masters NB, Eisenberg MC, Delamater PL, *et al*. Fine-Scale spatial clustering of measles nonvaccination that increases outbreak potential is obscured by aggregated reporting data. *Proc Natl Acad Sci U S A* 2020;117:28506–14.
- 18 Oyo-lta A, Wiysonge CS, Oranganje C, *et al*. Interventions for improving coverage of childhood immunisation in low- and middle-income countries. *Cochrane Database Syst Rev* 2016;7:Cd008145.
- 19 Kaboré L, Meda B, Médah I, *et al*. Assessment of missed opportunities for vaccination (Mov) in Burkina Faso using the world Health organization's revised Mov strategy: findings and strategic considerations to improve routine childhood immunization coverage. *Vaccine* 2020;38:7603–11.
- 20 Abbas K, Procter SR, van Zandvoort K, *et al*. Routine childhood immunisation during the COVID-19 pandemic in Africa: a benefit-risk analysis of health benefits versus excess risk of SARS-CoV-2 infection. *Lancet Glob Health* 2020;8:e1264–72.
- 21 Chandir S, Siddiqi DA, Mehmood M, *et al*. Impact of COVID-19 pandemic response on uptake of routine immunizations in Sindh, Pakistan: an analysis of provincial electronic immunization registry data. *Vaccine* 2020;38:7146–55.
- 22 World Health Organization. Pulse survey on continuity of essential health services during the COVID-19 pandemic. Available: [https://www.who.int/publications/i/item/WHO-2019-nCoV-EHS\\_continuity-survey-2020.1](https://www.who.int/publications/i/item/WHO-2019-nCoV-EHS_continuity-survey-2020.1) [Accessed 27 Aug 2020].
- 23 Tediosi F, Lönnroth K, Pablos-Méndez A, *et al*. Build back stronger universal health coverage systems after the COVID-19 pandemic: the need for better governance and linkage with universal social protection. *BMJ Glob Health* 2020;5:e004020.
- 24 Blanchard J, Washington R, Becker M. Vision 2035: public health surveillance in India. A white paper. NITI Aayog, government of India, 2020. Available: <https://niti.gov.in/node/1395>