


MSF Paediatric Days: a step forward in operationalising 'Humanitarian Paediatrics'

Sophie Janet ¹, Neal Russell,² Nikola Morton,³ Daniel Martinez,⁴ Mona Tamannai,¹ Nadia Lafferty,¹ Harriet Roggeveen,² Oluwakemi F. Ogundipe,⁵ Inmaculada Carreras,³ Anja Gao,³ Elise Didier,⁵ Roberta Petrucci⁴

To cite: Janet S, Russell N, Morton N, *et al.* MSF Paediatric Days: a step forward in operationalising 'Humanitarian Paediatrics'. *BMJ Paediatrics Open* 2021;**5**:e001156. doi:10.1136/bmjpo-2021-001156

Received 25 June 2021
Accepted 30 August 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Medical Department, Médecins Sans Frontières, Barcelona, Spain

²Medical Department, Médecins Sans Frontières, Amsterdam, The Netherlands

³Medical Department, Médecins Sans Frontières, Paris, France

⁴Medical Department, Médecins Sans Frontières, Geneva, Switzerland

⁵Medical Department, Médecins Sans Frontières, Brussels, Belgium

Correspondence to

Dr Sophie Janet; sophie.janet@barcelona.msf.org

ABSTRACT

Around the world, one in four children live in a country affected by conflict, political insecurity and disaster. Healthcare in humanitarian and fragile settings is challenging and complex to provide, particularly for children. Furthermore, there is a distinct lack of medical literature from humanitarian settings to guide best practice in such specific and resource-limited contexts. In light of these challenges, Médecins Sans Frontières (MSF), an international medical humanitarian organisation, created the MSF Paediatric Days with the aim of uniting field staff, policymakers and academia to exchange ideas, align efforts, inspire and share frontline research and experiences to advance humanitarian paediatric and neonatal care. This 2-day event takes place regularly since 2016. The fourth edition of the MSF Paediatric Days in April 2021 covered five main topics: essential newborn care, community-based models of care, paediatric tuberculosis, antimicrobial resistance in neonatal and paediatric care and the collateral damage of COVID-19 on child health. In addition, eight virtual stands from internal MSF initiatives and external MSF collaborating partners were available, and 49 poster communications and five inspiring short talks referred to as 'PAEDTalks' were presented. In conclusion, the MSF Paediatric Days serves as a unique forum to advance knowledge on humanitarian paediatrics and creates opportunities for individual and collective learning, as well as networking spaces for interaction and exchange of ideas.

INTRODUCTION

Around the world, one in four children are living in a country affected by conflict, political insecurity and disaster, and an unprecedented 30–34 million children are displaced from home.¹ Children are disproportionately impacted by crises and face the threat of violence, hunger, disease, disability and death. Worldwide, 56 million children under the age of 5 (half of them newborns) are projected to die between 2018 and 2030 in the absence of additional action, with the greatest proportion of this mortality anticipated in humanitarian and fragile settings.² Tragically, with a historic reversal of progress expected due

Key messages

What is known about the subject?

- ▶ One in four children worldwide are living in a humanitarian or fragile setting.
- ▶ There is a distinct lack of medical literature from humanitarian settings to provide guidance on best practice in such specific and resource-limited contexts.
- ▶ Médecins Sans Frontières (MSF) Paediatric Days was born to address paediatric issues of direct humanitarian concern.

What this review adds?

- ▶ This event unites frontline staff, policymakers and academia to exchange ideas, align efforts, inspire and share frontline research and experiences.
- ▶ Essential newborn care, community-based models of care, paediatric tuberculosis, antimicrobial resistance and collateral damage of COVID-19 on child health were discussed at the 2021 event.
- ▶ MSF Paediatric Days is as a unique forum to advance knowledge on humanitarian paediatrics and creates opportunities for individual and collective learning.

to the collateral impact of the SARS-CoV-2 pandemic, a rising gap between humanitarian needs and funding³ and declining support for child health in many cases, the number of children dying is expected to increase even further.⁴ In addition to increased mortality, we can expect escalating suffering, lost future potential and increasing inequality.^{5–7}

Médecins Sans Frontières (Doctors Without Borders/MSF) is an international medical humanitarian organisation specialising in responding to humanitarian emergencies such as conflicts, natural disasters and epidemics, acting with independence, neutrality and impartiality. MSF provides essential medical care to millions of children every year who would otherwise be without access to healthcare (see [table 1](#)).⁸

Healthcare in humanitarian and fragile contexts is challenging and complex to

Table 1 MSF paediatric activities involving children under 5 years old in 2019⁸

Description	n (rounded to closest whole)
Projects providing paediatric care	270
Children receiving outpatient care	3.3 million
Children receiving inpatient care unrelated to severe malnutrition	361 000
Children admitted to therapeutic feeding centres	186 000
Births supported	300 000

MSF, Médecins Sans Frontières.

provide, particularly for children. Healthcare staff are scarce, under-resourced and work well over capacity in some of the most insecure and adverse settings. Yet they continue to provide essential and life-saving services to vulnerable populations, rarely receiving the recognition they deserve. In addition, there is a distinct lack of medical literature from humanitarian settings to provide guidance on best practices in such specific and resource-limited contexts. There is a need to further integrate evidence-based practices into humanitarian contexts, and to generate evidence on what works best. We need to shine a brighter light on the experiences, challenges, failures and successes of those working in ‘humanitarian paediatrics’ (box 1) in order to improve care for this growing population of children in the most vulnerable circumstances.


There are not many platforms or venues available to bridge the existing gaps in clinical research and medical literature applicable to paediatric and neonatal care in humanitarian and fragile settings. Therefore, in 2016, the first *MSF Paediatric Days* was born with the aim of addressing urgent paediatric issues of direct humanitarian concern. Since then, three more editions have taken place at 18-month intervals, successfully uniting

Box 1 Humanitarian paediatrics

Humanitarian paediatrics refers to the branch of medicine that involves the care of newborns, infants, children and adolescents in humanitarian settings. It takes elements from public and global health, paediatrics, tropical and disaster medicine. It is centred on the child and his family and integrates the specific challenges and barriers related to the humanitarian context and limited resources.

frontline staff working in MSF and other organisations with policymakers and academia to exchange ideas, align efforts, inspire and share pertinent research and experiences. This event aims to raise awareness and exchange experiences, and to impact daily paediatric medical activities in humanitarian settings by promoting multidisciplinary collaboration, disseminating relevant best practices and generating specific recommendations (figure 1).

In April 2021, the first virtual edition of the *MSF Paediatric Days* brought together 1108 people from 95 different countries. MSF staff made up 58% of the attendees, largely frontline health workers, and the remainder came from a range of different organisations including academia, non-governmental organisations, ministries of health and other actors. The event included five main plenary discussions around key topics on humanitarian paediatrics, and eight virtual stands from internal MSF initiatives (MSF eCARE: electronic Decision Support System for paediatric primary care, Telemedicine: online tool for MSF medical frontline staff that provides access to specialised medical advice, point-of-care ultrasound) and external MSF collaborating partners (OPENPaediatrics, Save the Children, WHO, Laerdal and the American Academy of Pediatrics, Action Against Hunger, Alliance for International Medical Action and the Société Sénégalaise de Pédiatrie) who shared content and interacted with the participants. Additionally, 49 posters (available on ResearchGate) and 21 video abstracts were displayed,

**Figure 1** The MSF Paediatric Days approach to improving paediatric care in humanitarian settings. MSF, Médecins Sans Frontières.

offering evidence directly from hospitals and medical projects in fragile and humanitarian settings. This event provided a unique opportunity for frontline health staff working in such settings to share their experiences, challenges and solutions, in addition to creating networking spaces for interaction and exchange of ideas. Moreover, five inspiring short 'PAEDTalks' enriched the content of the event.

During this 2-day event on 15–16 April, five major themes were discussed: essential newborn care, community-based models of care, paediatric tuberculosis (TB), antimicrobial resistance in neonatal and paediatric care, and the collateral damage of the SARS-CoV-2 pandemic on child health. Replays and full event content can be found on the MSF Paediatric Days web page.

NEWBORN CARE: BACK TO BASICS

Newborns are one of the most vulnerable groups in humanitarian and fragile settings, but they have received limited focus in humanitarian action.⁹ The first session entitled 'Neonates - back to basics' illustrated the challenges that field teams face to implement, maintain, support and promote essential newborn care, with a focus on breastfeeding. Breastfeeding has well-recognised health benefits for mothers and newborns, but is even more crucial in humanitarian settings where breast milk substitutes are particularly dangerous in the absence of adequate water, sanitation and hygiene.^{10 11} Nevertheless, the life-saving nature of breastfeeding has often been overlooked in humanitarian settings, with a lack of institutionalised guidance on supporting breastfeeding.¹² The session identified numerous barriers to successful breastfeeding, such as false assumptions of breastfeeding being easy without need for support, contradictory messaging and failure to maintain mothers and babies together (mother–baby dyad care) due to lack of adequate space, resources and staff knowledge and awareness. Moreover, gender inequity and female disempowerment, which are exacerbated in crises, also negatively influence breastfeeding practices in many of the contexts where MSF intervenes.

There was a common recognition of the need for breastfeeding promotion and awareness raising among health staff, mothers and communities, including the need for professionals with expertise in breastfeeding promotion to support humanitarian responses. Ultimately, breastfeeding support needs to be recognised as an emergency humanitarian intervention. Different solutions and ideas were shared during the panel discussion to overcome some of these current challenges and to achieve this important shift in paradigm. Improving knowledge and skills on breastfeeding for healthcare providers is crucial, and support from lactation specialist should be made available via innovative platforms such as telemedicine. Engagement of all members of the community including traditional birth attendants, family (including male members) and any other key member

of the community should be included in the community engagement strategy of the project. Key messages of this session can be found in [table 2](#).

COMMUNITY-BASED MODELS OF CARE

The second session of the first day entitled 'Community-based models of care for neonatal and child health' provided insight on the current challenges and achievements of the decentralisation of care, and the important role that the community plays in the continuum of care. During the session, discussions highlighted the opportunity and importance of expanding community models of care for both children and newborns. Community health workers are crucial for delivering a range of preventive and curative health services, and for reducing inequities in access to care. It was highlighted that community activities should be built on existing capacity to avoid the implementation of parallel systems.

The integration of community-based models of care in emergency response is possible and most effective if the model is implemented in advance with standardised emergency preparedness strategies according to context, to promote resilience. Field testimonies from an Integrated Community Case Management programme in Niger and community-based nutrition programmes among others reinforced the importance, and the potential, that community models of care have in improving access and the continuum of care in children. See key messages of this session in [table 3](#).

PAEDIATRIC TB

On the second day, three main topics were discussed. The first session, 'Paediatric tuberculosis', touched on the challenges of diagnosing TB, the specificities of paediatric TB and the role of contact tracing and preventive treatment.

TB is a major infectious killer in MSF settings and children, especially those under 5 years of age, are at particular risk of severe forms of the disease. Paediatric TB is a 'silent disease' frequently underdiagnosed, undertreated and under-reported. TB presents differently in children than in adults, with a higher proportion of extrapulmonary TB. Microbiological confirmation is rarely achievable in children and is especially challenging in humanitarian settings. Therefore, emphasis on clinical diagnosis is imperative to ensure that presumptive treatment is started without delay. Treatment of paediatric TB based on clinical diagnosis alone would decrease TB morbidity in children, thereby minimising preventable deaths from the disease.

Contact tracing of TB cases in the community and subsequent tuberculosis preventive treatment (TPT) is often overlooked and deprioritised in low-resource settings, but this should be pursued more actively. This can now be facilitated with new, shorter TPT regimens, which have already shown promising results in terms of

**Table 2** Newborn care: back to basics

Key messages	Why is it important?	Current challenges	Recommendations
Breast feeding (BF) is an intervention that saves lives, improves health and development of newborns, as well as maternal well-being. BF should be universally and practically achieved with dedicated support in all MSF contexts.	Newborn mortality and morbidities remain high across MSF projects. Essential evidence-based interventions shown to decrease newborn mortality such as exclusive and early BF should be supported and scaled up to save lives across MSF. BF is natural, instinctive, ready made and vastly available. However, many women face different challenges to establish and sustain BF. To overcome those challenges, a coordinated and multidisciplinary support should be available for every woman and their baby.	<ul style="list-style-type: none"> ▶ BF is believed to be intuitive and easy for women. This is globally recognised as a harmful assumption. ▶ Essential, evidence-based aspects of BF, such as to starting within the first hour of life and exclusive BF for 6 months, are not always considered. ▶ BF is not always recognised as an intervention and therefore there are no allocated resources for BF support. ▶ Suboptimal training and preparation lead to varying and even contradictory messages given to the mother and family within MSF projects. 	<p>Field</p> <ul style="list-style-type: none"> ▶ Consider BF as an intervention to reduce newborn mortality and allocate space, time and resources in planning for it. ▶ Promote BF and essential newborn care champions or focal points. ▶ Support and promote early and exclusive BF, including where it seems not easy for newborn or mother. ▶ Promote multidisciplinary (midwife, nutritionist, nurses, doctors, logistician) work to support BF, increase awareness and discuss responsibilities and division of tasks. <p>Operations</p> <ul style="list-style-type: none"> ▶ Include essential newborn care interventions (such as BF and Kangaroo Mother Care) into the main/strategic interventions to decrease neonatal mortality at project level and coordinate resources to support it. ▶ Promote partnership with other actors involved in essential newborn care, especially at local level. <p>Headquarters</p> <ul style="list-style-type: none"> ▶ Ensure BF policies and guidance are available and harmonised across MSF. ▶ Support and encourage access to lactation consultants in telemedicine or other platforms to support field teams. ▶ Ensure that training on essential newborn care including BF is available in different languages for frontline staff.
A family-centred approach, which includes an understanding of the community and the context, is needed to ensure successful BF.	To effectively support mothers, we need to understand the barriers and enablers related to a specific context. The mother–baby dyad is at the centre of the process, but all the family and community need to participate, support, encourage.	<ul style="list-style-type: none"> ▶ There is often little understanding about how BF is perceived in different contexts and what are the barriers and enablers in different settings, including the influence of other family members. 	<p>Field</p> <ul style="list-style-type: none"> ▶ Include families, caretakers, community health workers in understanding the local influences to support the promotion of BF. <p>Operations</p> <ul style="list-style-type: none"> ▶ Essential newborn care (including BF understanding and support) should be factored into community-level programmes. <p>Research</p> <ul style="list-style-type: none"> ▶ If BF levels are low or poorly understood, consider anthropological studies in different contexts on barriers and enablers for BF. Include male views.

BF, Breastfeeding; MSF, Médecins Sans Frontières.

acceptance, efficacy, safety and adherence to treatment within MSF settings. See key messages of the session in [table 4](#).

ANTIMICROBIAL RESISTANCE AND ANTIMICROBIAL STEWARDSHIP IN NEONATAL AND PAEDIATRIC CARE

The second topic discussed on 16 April was ‘Antimicrobial resistance and antimicrobial stewardship in neonatal and paediatric care’. Antimicrobial resistance is a reality in humanitarian settings^{13 14} and has been described as a silent tsunami. Vulnerable groups such as newborns and malnourished children face a disproportionate burden

and specific challenges. The challenges and consequences of outbreaks in neonatal units in low-resource settings were highlighted, including both experience within MSF and growing evidence based in low-resource settings outside MSF.^{15 16} In addition, MSF field experiences from Mali and South Sudan underlined the need for a multidisciplinary approach focusing on several transversal pillars including infection prevention and control (IPC), antibiotic stewardship and microbiology (when possible or available). Antibiotic stewardship is particularly challenging for paediatric patients in humanitarian settings. The combination of high mortality and a lack

Table 3 Community-based models of care in paediatrics

Key messages	Why is it important?	Current challenges	Recommendations
Community models of care are effective in delivering a range of preventive, promotive and curative health services for children and neonates, and they can contribute to reducing inequities in access to care.	<p>In humanitarian and fragile settings when access to health facilities is limited, care at community level can bridge important health gaps for mothers, newborns and children.</p> <p>The community-based activities are an essential part of the health system, contributing to build skills and confidence to empower people with knowledge, tools and understanding referral needs.</p>	<ul style="list-style-type: none"> ▶ Community activities suffer from lack of anchorage with the existing health system and tend to be implemented as a parallel system. ▶ Monitoring and evaluation (M&E) of the service delivered is hampered by the lack of clear and simple core indicators. ▶ Community health workers (CHW) are given more and more responsibilities, their skills and workload not always match. 	<p>Field</p> <ul style="list-style-type: none"> ▶ Community models of care should be rooted in understanding of the context, social realities and values of the communities we are working with and designed in a participatory manner. ▶ Simplified core indicators of implementation, quality of care and utilisation of services should be implemented to allow M&E, along with qualitative data to understand important barriers and enablers. ▶ Involve communities in M&E of programmes at a minimum through assuring context-appropriate feedback mechanisms are in place. ▶ Ensure realistic workload of the CHW and enhance their motivation through social recognition of their work, an appropriate reward system, regular supervision, feedback, exchanges, sense of belonging to a larger network. <p>Operations</p> <ul style="list-style-type: none"> ▶ Community activities should be built on existing capacity, avoiding the implementation of a parallel system. <p>Headquarters</p> <ul style="list-style-type: none"> ▶ Provide a framework for assessing/training CHWs and a catalogue of relevant expectations of CHW dependant on achievable and most relevant competencies.
Community models of care in emergency response are most effective if the model is implemented in advance with contextual emergency preparedness (EPREP) strategies.	Empowering the community in delivering healthcare increases resilience during crises when access to the health facilities may be further limited.	<ul style="list-style-type: none"> ▶ Planning and preparation are essential to deliver effective emergency response, but there is still little investment in EPREP at community level. 	<p>Field/operations</p> <ul style="list-style-type: none"> ▶ Integrate paediatric and neonatal community activities in the EPREP strategy. <p>HQ</p> <ul style="list-style-type: none"> ▶ Further simplify tools, M&E indicators and a framework for prioritisation for community activities during emergency response.

CHW, Community health workers; EPREP, Emergency preparedness; HQ, Headquarters; M&E, Monitoring and Evaluation.

of microbiology testing means that severe illness is often treated with diagnostic uncertainty, leading to an overuse of antibiotics. The challenge of access to microbiology was discussed, including the important role that understanding of local antimicrobial resistance patterns plays in stewardship practices. While extensive data on antimicrobial resistance exist in some settings, there is a complete lack of data in many parts of the world. But even without access to microbiology, there is capacity to improve antibiotic stewardship and IPC, which are essential and feasible in all settings.¹⁷ The importance of an interdisciplinary approach was discussed, ensuring involvement of all members of the team, including doctors, nurses, pharmacists and cleaners. Practical tools to support field teams to assess and monitor medical activities from an antimicrobial resistance lens are available within and

outside MSF, such as point prevalence surveys on antibiotic use, antibiotic consumption analysis and the stepwise IPC approach. Key messages for this session can be found in [table 5](#).

COLLATERAL DAMAGE OF COVID-19 ON CHILD HEALTH

The last session of the event was dedicated to the ‘Collateral damage of COVID-19 on child health’. Children have been disproportionately affected by the pandemic, with low mortality due to COVID-19 itself, but high morbidity and mortality due to the multiple collateral effects of the health crisis. The pandemic has impacted child health through increases in poverty, loss of education, food insecurity and violence, as well as greater strains on health systems and a reduction in access to

Table 4 Paediatric tuberculosis

Key messages	Why is it important?	Current challenges	Recommendations
Underdiagnosis and undertreatment of paediatric tuberculosis (TB) lead to preventable deaths. Microbiological confirmation is rarely available in children, therefore at present, a clinical diagnosis should be used to start presumptive treatment without delay.	<p>TB remains a major, unrecognised killer in children. MSF has a possibility to make a difference now by increasing the knowledge of field teams who meet children or their caretakers.</p> <p>Presumptive and empirical TB treatment is safe, well tolerated and effective. Starting treatment based on clinical suspicion (not microbiology confirmation) will bridge the gap of underdiagnosis and undertreatment of TB in children in MSF projects.</p>	<ul style="list-style-type: none"> ▶ Paediatric TB is a 'silent disease' frequently underdiagnosed, undertreated and under-reported. ▶ MSF staff are not always familiar with the different clinical presentations of TB in children and there is a gap in capacity building on this topic. ▶ Confirmatory TB diagnosis is often hard to access and can be difficult in children. ▶ Delays of starting treatment based on a microbiological diagnosis perpetuate TB undertreatment in children who may die through these unnecessary delays. 	<p>Field</p> <ul style="list-style-type: none"> ▶ Know the local burden of paediatric TB. ▶ Support medical field teams on how to recognise TB in children as part of their daily work. ▶ While caring for adults with TB, consider the children exposed. ▶ Treat TB based on clinical suspicion. <p>Operations</p> <ul style="list-style-type: none"> ▶ Integrate TB activities in paediatric care. ▶ Monitor programme data and investigate if underdiagnosis is suspected depending on the local prevalence of TB. ▶ Promote capacity building and facilitate access to learning opportunities on paediatric TB including the online free course. <p>Headquarters/research</p> <ul style="list-style-type: none"> ▶ Advocate for the integration of TB in all paediatric projects. ▶ Provide support and guidance on clinical algorithm for the diagnosis and treatment of TB in paediatric projects.
Tracing the contacts of patients with TB with the offer of tuberculosis preventive treatment (TPT) should be pursued as an effective strategy to save lives in MSF projects.	<p>Contact tracing of patients with TB is an effective way to identify those who have active TB but also those who may be harbouring latent (sleeping) TB. More lives can be saved by improving access to timely treatment or TPT.</p> <p>New shorter drug regimens for TPT are showing promising results on acceptance, effectivity, safety and adherence to treatment.</p>	<ul style="list-style-type: none"> ▶ Contact tracing requires resources, which is often a barrier to its roll-out in communities, especially if it is in addition to other community activities. ▶ Standard TPT strategy is currently well established, but shortens regimens that show promising results have not been fully validated for MSF programmes. 	<p>Field</p> <ul style="list-style-type: none"> ▶ Contact tracing should be performed whenever a TB case is identified. ▶ Assure systematic follow-up of children under TPT in the community. <p>Operations</p> <ul style="list-style-type: none"> ▶ Innovate and pilot TPT programmes in settings where the need is clear and share experiences with the whole MSF movement to improve future efforts. ▶ Seek partnership for TPT with community and other non-governmental organisations to reduce the resource burden and optimise programme reach. <p>HQ/working groups</p> <ul style="list-style-type: none"> ▶ Determine where TPT will be most beneficial to reduce paediatric TB burden and implement and learn from those MSF sites.

HQ, Headquarters; MSF, Médecins Sans Frontières; TB, Tuberculosis; TPT, Tuberculosis preventive treatment.

health services. Preventive services like vaccination and nutrition programmes have been mostly suspended or delayed.¹⁸ The indirect effects of the pandemic have been most severe in resource-limited settings where increased child mortality is a major concern, widening the gap of inequity for children. There were field testimonies from frontline health workers outlining the struggle they faced during this unprecedented time. Supporting

health systems to maintain preventive and curative services is crucial to attenuate the ongoing impact. Flexibility to adjust health activities is key to face the challenges brought by the SARS-CoV-2 pandemic. Boosting community healthcare activities within MSF strategies, as an essential part of the continuum of care, is an efficient way of assuring healthcare access. Find in [table 6](#) the key messages of this session.

Table 5 Antimicrobial resistance and antimicrobial stewardship in neonatal and paediatric care

Key messages	Why is it important?	Current challenges	Recommendations
<p>Patients, and especially newborns and children, are harmed by and even die because of antimicrobial resistance (AMR) in MSF projects. The problem is escalating in front of us like an invisible tsunami, with limited visibility on its burden and consequences.</p> <p>It is critical for MSF to systematically implement the available tools to reduce AMR, especially where microbiology is not available: infection prevention and control (IPC) and antibiotic stewardship</p>	<p>AMR is a reality in humanitarian settings and newborn and children are particularly exposed.</p> <p>Multidrug-resistant bacterial sepsis particularly affects the most fragile patients, as shown by the increase in the reports of outbreaks in neonatal units in low-resource settings.</p> <p>IPC and antibiotic stewardship are crucial and effective strategies against AMR, particularly in contexts where microbiology is unavailable.</p>	<ul style="list-style-type: none"> ▶ There is lack of awareness on the increasing paediatric and neonatal morbidity and mortality because of AMR in humanitarian settings. ▶ There is a false perception that AMR does not affect low-resource settings and limited available data to accurately define the extent of the problem. ▶ Misconception that without microbiology, it is not possible to tackle AMR. ▶ There are gaps on access to microbiological tools. 	<p>Field</p> <ul style="list-style-type: none"> ▶ Strengthen awareness and training on IPC interventions, and scale up use of IPC quality improvement tools. ▶ Create multidisciplinary AMR project committee including all the relevant health workers (nurses, doctors, pharmacists, IPC focal points, cleaners), and identify focal points and champions. ▶ Scale up use of audits of antimicrobial use. <p>Operations</p> <ul style="list-style-type: none"> ▶ Integrate AMR and antibiotic stewardship as part of quality improvement initiatives. ▶ Formalise AMR and IPC focal point roles in job descriptions. ▶ Increase access to microbiological tools available to the field, including exploring partnerships with national and regional laboratories. <p>Research/Headquarters</p> <ul style="list-style-type: none"> ▶ Adapt IPC assessment tools to address specific challenges in neonatal and paediatric care. ▶ Update guidelines in accordance with evidence on AMR in different infection syndromes. ▶ Explore alternative metrics/indicators for antibiotic use in children to guide antibiotic stewardship.

AMR, Antimicrobial resistance; IPC, Infection prevention and control; MSF, Médecins Sans Frontières.

Table 6 Collateral damage of COVID-19 on child health

Key messages	Why is it important?	Current challenges	Recommendations
<p>Children have disproportionately been affected by the COVID-19 pandemic, with low direct mortality, but high morbidity and mortality due to the multiple collateral effects of the health crisis.</p> <p>This unprecedented crisis offers an opportunity to change our ways of thinking, deploy and maintain our activities, rethink support models and define future preparedness and responses.</p>	<p>The pandemic has impacted child health through increases in poverty, loss of education, food insecurity, violence as well as increased strain on health systems and reduction in access to health services.</p> <p>These collateral effects of the pandemic have been most striking in resource-limited settings where increase in child mortality is a major concern.</p> <p>Preventive services like vaccination and nutrition programmes have been affected most by suspension or delay.</p> <p>The tremendous detrimental effects of the pandemic on child health are still unfolding and our concern as MSF should be high.</p>	<ul style="list-style-type: none"> ▶ The focus on the direct impact of COVID-19 has had a huge and overlooked negative impact on children through the reduction and suspension of essential healthcare services. ▶ The risk of weakening essential services continues through resurgence of the pandemic. There is a potential extra burden related to vaccination programmes in poorly resourced health systems, where healthcare workers and resources will be repurposed to deliver those vaccines at the expense of critical childhood services. 	<p>Field</p> <ul style="list-style-type: none"> ▶ Adapt and innovate to maintain routine services, such as continuum of nutritional screening and vaccination. ▶ Witness, document and report collateral effects of the pandemic on children, real time and in retrospect. <p>Operations</p> <ul style="list-style-type: none"> ▶ Maintain the preventive and curative paediatric regular services to limit an increase in child morbidity and mortality. ▶ Boost the community healthcare activities in MSF strategies as an essential piece of the continuum of care and as an efficient way of assuring health access. ▶ Be flexible and innovative in order to adjust our healthcare activities according to the situation and to provide proper technical medical support to the field teams. <p>Headquarters</p> <ul style="list-style-type: none"> ▶ Consider COVID-19 pandemic as a transformative opportunity to develop new approaches and implement new and practical tools needed in the field reality. ▶ Advocate at national and international levels for the continuity of routine preventive and curative paediatric and neonatal activities in this pandemic.

MSF, Médecins Sans Frontières.

Despite the collateral damage already caused, this past year can also be an opportunity to change our ways of thinking, activities, support models and future preparedness and responses. More than ever, now is a crucial time to invest in humanitarian paediatrics, to support children and uphold their rights in the most fragile contexts.

CONCLUSIONS

As the world continues to battle the SARS-CoV-2 pandemic, focus and funding for child health in humanitarian settings suffer while children's needs escalate. The future for children in humanitarian settings hangs in the balance, and platforms that raise the profile of humanitarian paediatrics are vital to ensure that these children are not overlooked and remain a priority among funders, decision-makers and stakeholders. The MSF Paediatric Days serves as a unique forum to advance knowledge on humanitarian paediatrics, and to create opportunities for individual and collective learning on this topic. We look forward to the next edition and welcome all suggestions for the next topics of the MSF Paediatric Days.

Acknowledgements We are grateful to our frontline field staff for providing essential and life-saving services to vulnerable populations, despite the enormous challenges posed by the contexts in which they work. We would like to acknowledge their commitment to their patients and the incredible work that they carry out on a daily basis to decrease the inequalities in child health globally.

Contributors SJ: substantial contributions to the conception, drafting of manuscript, revision, final approval and submission. NR, NM, DM, MT, NL, OO, ED, RP: substantial contributions to the conception, revision and final approval. HR, IC, AG: substantial contributions to the conception and final approval.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Commissioned; externally peer reviewed.

Data availability statement Data sharing not applicable as no data sets generated and/or analysed for this study.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Sophie Janet <http://orcid.org/0000-0003-0194-2406>

REFERENCES

- 1 United Nations Office for the Coordination of Humanitarian Affairs. Global humanitarian overview 2019 [online], 2019. Available: <https://reliefweb.int/sites/reliefweb.int/files/resources/GHO2019.pdf>
- 2 UNICEF. *Levels and trends in child mortality report 2018*, 2018.
- 3 A.Spencer BW-K&. *Reducing the humanitarian financing gap: review of progress since the report of the high-level panel on humanitarian financing*, 2021.
- 4 Robertson T, Carter ED, Chou VB, *et al*. Early estimates of the indirect effects of the COVID-19 pandemic on maternal and child mortality in low-income and middle-income countries: a modelling study. *Lancet Glob Health* 2020;8:e901–8.
- 5 Wagner Z, Heft-Neal S, Bhutta ZA, *et al*. Armed conflict and child mortality in Africa: a geospatial analysis. *Lancet* 2018;392:857–65.
- 6 Shenoda S, Kadir A, Pitterman S, *et al*. The effects of armed conflict on children. *Pediatrics* 2018;142:e20182585.
- 7 Pfefferbaum B, Jacobs AK, Van Horn RL, *et al*. Effects of displacement in children exposed to disasters. *Curr Psychiatry Rep* 2016;18:1–5.
- 8 MSF Paediatric Working Group. *Paediatric activities medecins SANS Frontieres*, 2019.
- 9 UNICEF. *Save the children WHO. roadmap to accelerate progress for every newborn in humanitarian settings 2020–2025*, 2020.
- 10 Jakobsen M, Sodemann M, Nylén G, *et al*. Breastfeeding status as a predictor of mortality among refugee children in an emergency situation in Guinea-Bissau. *Trop Med Int Heal* 2003;8:992–6.



- 11 Hipgrave DB, Assefa F, Winoto A, *et al.* Donated breast milk substitutes and incidence of diarrhoea among infants and young children after the may 2006 earthquake in Yogyakarta and central Java. *Public Health Nutr* 2012;15:307–15.
- 12 Shaker-Berbari L, Ghattas H, Symon AG, *et al.* Infant and young child feeding in emergencies: organisational policies and activities during the refugee crisis in Lebanon. *Matern Child Nutr* 2018;14:e12576.
- 13 Kanopathipillai R, Malou N, Hopman J, *et al.* Antibiotic resistance in conflict settings: lessons learned in the Middle East. *JAC Antimicrob Resist* 2019;1:2–4.
- 14 Langendorf C, Le Hello S, Moumouni A, *et al.* Enteric bacterial pathogens in children with diarrhea in niger: diversity and antimicrobial resistance. *PLoS One* 2015;10:e0120275.
- 15 Okomo U, Senghore M, Darboe S, *et al.* Investigation of sequential outbreaks of Burkholderia cepacia and multidrug-resistant extended spectrum β -lactamase producing Klebsiella species in a West African tertiary hospital neonatal unit: a retrospective genomic analysis. *The Lancet Microbe* 2020;1:e119–29.
- 16 Lenglet A, Faniyan O, Hopman J. A nosocomial outbreak of clinical sepsis in a neonatal care unit (NCU) in Port-au-Prince Haiti, July 2014 – September 2015. *PLoS Curr* 2018.
- 17 Johnson J, Akinboyo IC, Curless MS, *et al.* Saving neonatal lives by improving infection prevention in low-resource units: tools are needed. *J Glob Health* 2019;9:9–11.
- 18 Sidhu S. *WHO and UNICEF warn of a decline in vaccinations during COVID-19.* World Health Organization, 2020.