298

A COMPARATIVE STUDY OF CLINICAL PROFILE AND SYMPTOM CONTROL IN OVERWEIGHT AND NORMAL WEIGHT SCHOOL AGE CHILDREN WITH MILD PERSISTENT ASTHMA

Sharmila Manivannan, Venkatesh Chandrasekaran, Nandheeswari Subramanian. *UK*

10.1136/bmjpo-2021-RCPC.166

Background Bronchial asthma and obesity show a parallel increase in prevalence of late. Obese children with asthma have been shown to have lower disease control and increased symptom severity compared to normal weight asthmatics. There has been a dearth of studies related to asthma in obese and overweight children in India.

Objectives To find out the proportion of children with poor symptom control in overweight/obese and normal weight children with mild persistent asthma and to know the socio-demographic and clinical correlates of poor symptom control in them.

Methods Children aged 6 to 12 years with mild persistent asthma with BMI Z score for age and sex more than +1 Z score on WHO BMI Z score chart for age and sex formed the cases. Age and sex matched asthmatics with BMI Z score for age and sex between -2 Z to +1 Z score formed the controls. A brief, relevant history and respiratory examination was carried out in them. FEV1, FEV1/FVC were measured in both groups using Care Fusion Jaeger spirometer. Symptom control was assessed by ACT score and we also looked at inhaler device technique. Statistical analysis was done using SPSS version 19 and Vassarstats.

Results Our study population consisted of predominantly males hailing from rural areas. Parental atopy was found in equal proportions among both cases and controls. All most all children in the overweight/obese group showed good compliance to therapy. The proportion of children with poor control was 19.1% in the overweight/obese group and 23.4% in the normal weight group. There was no significant correlation between BMI and symptom control as assessed by ACT score. The lung function tests performed in our study population revealed a mean FEV1 (% of predicted) of around 75% in both the groups in children with good control. This relates to the mild obstructive lung dysfunction in our study population. For those children with poor control, the mean FEV1 (% of predicted) was around 67% in both the groups. Overweight/obese children with good control showed a slightly lower FEV1/FVC ratio compared to children with normal weight. Overweight/obese children also had a slightly higher median eosinophil count compared to children with normal weight. Gastro-oesophageal reflux and allergic rhinitis were more commonly seen in overweight/obese children. In the poor control group, FEV1, FEV1/FVC and median eosinophil counts were not significantly different between overweight/obese and

Abstract 298 Table 1 Comparison of symptom control in the study groups

Symptom control	Overweight/ Obese (n=47)	Normal weight (n=47)	P value*
Good control (ACT $\geq$ 20)	38	36	0.25
Poor control (ACT<20)	9	11	0.065

*P value calculated using Chi square test

normal weight group, but was less when compared to good control group.

Conclusions The proportion of poor symptom control in overweight/obese children with mild persistent asthma is not significantly different when compared to normal weight children with mild persistent asthma as assessed by ACT score and lung function measurements. No significant risk factors for poor symptom control could be identified in both overweight/obese group as well as normal weight group in our study.

302

VIRTUAL PAEDIATRIC CLINIC IN SPECIAL NEEDS SCHOOL- IS THAT RIGHT FOR MY CHILD?

Tapomay Banerjee, Amjad Khan, Piriya Kesavan, Samira Ajmal, Rishi Arora. *UK*

10.1136/bmjpo-2021-RCPC.167

Background Community paediatric services at Bedford, UK offers clinical support to all 3 local special needs schools. A multidisciplinary clinic with a paediatrician and special needs nursing team takes place once a week, and with child and adolescent mental health services team once in 6–8 weeks at the school sites. However, because of COVID-19 pandemic all schools closed in March 2020 and we had to stop face to face clinics. We converted these clinics into initially telephone and subsequently either telephone/video consultations.

Objectives The objective of our study was to assess how parents of children and young people (CYP) with special needs feel about this transition to virtual clinics.

Methods CYP attending special schools in Bedford, who are under the care of community paediatricians (n=397), were selected for the study. Parents of children, who were seen in virtual clinics between 01/07/2020 and 31/10/2020, were approached for feedback (n=110). This was part of a wider survey related to these children's anxiety and emotional difficulties during lock-down. A link to the questionnaire was sent to parents' phone/Email after explaining the study and obtaining verbal consent. All responses were recorded in the IQVIA Connection survey website (<https://www.oc-meridian.com/cambsCommunityServices>). No sensitive or personal identifiable data were used.

Abstract 302 Table 1 Comments made by parents

Parental Experience of virtual clinics	<ol style="list-style-type: none"> 1. Doctor listens and gives good advice The centre is really helping and give a great follow up to parents of children with disabilities Be reassured that everything will be ok 2. Everyone has been extremely helpful during this situation. I cannot fault a single thing. 3. We discussed different elements of my daughters needs and agreed on a plan going forward, I felt included and that I mattered in the decision making process 4. Overall the Nursing Team and the Paediatrician have been very supportive over the lockdown. 5. I thought it was a great idea. I had to cancel at short notice - waiting for another time but we have no issues so I'm not chasing.
--	---

Results We received 53 responses from parents. 31(58.49%) have neurodevelopmental disorder (autism spectrum disorder/attention deficit hyperactivity disorder), 14(26.41%) have learning disabilities, 5(9.43%) have neuro disabilities (cerebral palsy/global developmental delay, and 3(5.66%) have other conditions such as chromosomal deletions.

27(50.94%) parents felt that a virtual appointment with a paediatrician and nurses were very good and another 16 (30.18%) felt it was good. 2(3.77%) of them felt it was neither good nor poor. None of them felt it was poor or very poor service but 8(15.09%) either did not answer this question or said I don't know. We also requested feedback from them about their experience and their comments are summarized in table 1.

Conclusions Overall, parents felt very much reassured that we are running a service during lockdown. However one parent felt that her child was too complex to be seen in virtual clinic. Our observations support the idea of multidisciplinary virtual clinics for special needs children but we agree that it cannot replace face to face consultations.

305 PROFILES OF MALTREATED CHILDREN AGED 0–3 YEARS OLD AND THEIR FAMILIES: THE SINGAPORE EXPERIENCE

Rui Kwan, Sharon Abraham, Wan Har Chong, Jean Yin Oh, Yeleswarapu Sita Padmini. Singapore

10.1136/bmjpo-2021-RCPCH.168

Background Early adverse childhood experiences (EACES) such as child maltreatment between the ages of 0 to 3, has significant impact on a child's cognitive development and behaviour. Negative experiences and even the absence of positive experiences e.g. in extreme deprivation, can cause disruptions to brain architecture, with long-lasting effects. EACES also increase the risk of physical health, mental health and relational difficulties in adulthood with poor academic and employment outcomes. Understanding the risk factors associated with health and developmental conditions in maltreated children provide more effective interventions to improve their outcomes.

Objectives The purpose of the study was two-fold. One, to explore the demographics, social, developmental and physical health conditions of young children admitted to a tertiary children's hospital for suspected child maltreatment Two, to examine the occurrence of child maltreatment with developmental delays and missed vaccinations.

Methods 99 children aged 0 to 3, admitted between the period of January 2018 and June 2019 were included in this quasi-observational study. Medical data was electronically extracted on variables surrounding their health, development, and demographics.

Results Children in this study had a mean age of 1.3 years; and 59% were males. The most common reported abuse type was physical (74%), followed by neglect (14%) and sexual abuse (10%) 2% had more than one of three abuse types. 35% of children had pre-existing developmental conditions and significant medical conditions e.g. asthma or atopy. During the hospitalisation, 58% of children were assessed to have child maltreatment-related injuries, missed vaccinations and developmental conditions requiring health care follow-up. Specifically, 19% ($n=19$) of children had existing or

present developmental delays, where six were detected through routine developmental screening during hospitalization. Children with developmental delays were more likely to have missed vaccinations than those without delays [Odds ratio (OR) 3.6, 95% confidence interval (CI) 1.04–12.62]. The correlation with developmental delays was independent of maternal education or household income. Children staying in rented housing were more likely to have missed vaccinations than those staying in purchased housing [OR 9.5, 95% CI 1.09–82.72]. When it came to parental factors, the majority of children's parents had only primary school education; and had a mean household income of S\$823 – well below the national average. 20% stayed in purchased housing, 27% in rented housing, and 5% were lodging with relatives or friends. Majority of children (72%) were referred to statutory child protection agencies as part of safeguarding procedures, indicating significant concerns over the child's injury and social circumstances.

Conclusions A sizeable proportion of maltreated children were identified with developmental delays – twice that seen in the general population. A significant proportion also presented with health concerns that utilize health care resources and follow-ups. Those with social deprivation were more likely to have neglected basic health needs e.g. vaccination. Thus, managing child maltreatment should include enhanced developmental screening using standardized tools to improve early detection and robust health surveillance, ensuring the timely identification of needs and access to appropriate services for better health equity.

306 RENAL VASCULAR LESIONS IN CHILDHOOD-ONSET LUPUS NEPHRITIS: A TERTIARY CENTRE EXPERIENCE

Eugene Yu-hin Chan, Alison Lap-tak Ma, Yuen-fun Mak, Wai-ming Lai. Hong Kong

10.1136/bmjpo-2021-RCPCH.169

Background The clinical significance of renal vascular lesions remains unknown in childhood-onset lupus nephritis (LN) and is not currently factored into WHO and ISN/RPS classification.

Objectives The aim of this study is to determine the incidence and outcomes of childhood-onset LN in the presence of these lesions.

Methods We conducted a retrospective review at the tertiary paediatric nephrology centre in Hong Kong of children with LN who presented before 18 years with active follow-ups for 12 or more months by July 2020. Estimated glomerular filtration rate (eGFR) was calculated by modified Schwartz and CKD-EPI formula as appropriate.

Results Among 90 children with childhood-onset systemic lupus erythematosus, 61 patients (68%; mean age of 13.2 ± 3.4 years; 85% girls) diagnosed to have biopsy-proven LN. The predominant presentation was nephritic-nephrotic syndrome ($n=30$, 50%), with 4 patients required acute dialysis. One (4%), 17 (28%), 29 (48%), 9 (15%), 4 (7%) and 1 (2%) patients had class II (lupus podocytopathy), III, IV, III/IV + V, V and VI LN, respectively.

Of the 61 patients with LN, renal vascular lesions were observed in 9 children (15%) at a mean age of 14.1 ± 4.3 years, including lupus vasculopathy (LV; $n=4$, 7%), thrombotic microangiopathy (TMA; $n=4$, 7%) and true vasculitis ($n=1$, 2%). Eight patients had co-existing proliferative LN (Class III