


# Epidemiological characteristics of paediatric burn patients in China from 2016 to 2019: a retrospective study

Kun Chen,<sup>1</sup> Jing Liu,<sup>1</sup> Xinyu Wang,<sup>2</sup> Ping Chu,<sup>3</sup> Guoshuang Feng,<sup>2</sup> Yueping Zeng,<sup>4</sup> Xin Xu,<sup>5</sup> Yongli Guo,<sup>3</sup> Xin Ni <sup>3,6</sup>

**To cite:** Chen K, Liu J, Wang X, *et al.* Epidemiological characteristics of paediatric burn patients in China from 2016 to 2019: a retrospective study. *BMJ Paediatrics Open* 2023;**7**:e001796. doi:10.1136/bmjpo-2022-001796

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjpo-2022-001796>).

KC and JL contributed equally.

Received 24 November 2022

Accepted 25 March 2023

## ABSTRACT

**Background** Epidemiological characteristics of paediatric burn can help to protect children from burn injuries. Most of the previous studies have been conducted on small scale and single centre in China. This study analysed the epidemiological characteristics of paediatric burn using a large-scale and multicentre database from 23 children's hospitals in China to increase child protection against burn injuries, improve the quality of care and reduce hospitalisation costs.

**Methods** Excerpted information from medical records of 6741 paediatric burn cases was extracted from the Futang Research Center of Pediatric Development database from 2016 to 2019. The epidemiological characteristics of patients, including gender, age, cause of burn injuries, complication, the timing of hospitalisation (season and month) and hospitalisation length and cost, were collected.

**Results** Male gender (63.23%), aged 1–2 years (69.95%), and hydrothermal scald (80.57%) were significantly dominant among cases. Furthermore, complications were significantly different among groups with different age of patients. Pneumonia was the most common complication (21%). Meanwhile, most paediatric burn occurred in spring (26.73%), and hospitalisation length and cost significantly depended on the cause of the burn injuries and surgical intervention.

**Conclusions** This large-scale epidemiological study of paediatric burn in China revealed that boys aged 1–2 years with higher activity and lack of self-awareness are more likely to suffer from burn injuries by hydrothermal scald. In addition, complications, especially pneumonia, need attention and early prevention in paediatric burn.

## BACKGROUND

Burn injuries are the major type of accidental injuries in children globally,<sup>1</sup> with heavy economic and psychological burdens for families and society.<sup>2,3</sup> The children suffering from burn injuries in Asia account for more than half of paediatric burn cases worldwide, most of whom require hospitalisation.<sup>4,5</sup> However, as a country with a huge number of children, China does not have adequate epidemiological information about the paediatric burn. Most of previous epidemiological reports on paediatric burn have been small-scale studies.<sup>6,7</sup> A large-scale and multicentre

## WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Burn injuries are the main type of accidental injuries in children, and most of the epidemiological studies on this topic in China are small scale and single centre.

## WHAT THIS STUDY ADDS

⇒ Scalds were the most frequent cause of burn injuries.  
⇒ Toddlers aged 1–2 years were the most frequently affected group.  
⇒ Pneumonia was the most frequent complication.

## HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Education programmes for burn prevention targeting families, schools and child-care facilities are an important public health initiative to reduce the burn burden. According to the epidemiological information of paediatric burns, medical centres can devote intensive resources to prepare each patient and provide the best service. More effective evidence-based, individualised programmes should be developed to treat paediatric burn with high complication rates.

study can provide much more accurate information about paediatric burn, which can be used to promote care quality in paediatric burn. Therefore, it is necessary to retrospectively analyse the epidemiological characteristics of paediatric burn in a large-scale and multicentre study.

The Futang Research Center of Pediatric Development (FRCPD) is a large medical consortium and a non-profit social service organisation in China, greatly promoting research in the paediatric field.<sup>8</sup> The FRCPD covers almost one-third of children's hospitals in China. In our prior studies, we reported that exposure to hot materials is the fifth cause of morbidity and mortality by analysing 122 284 medical records in the FRCPD database.<sup>9</sup> Using the FRCPD database in this study, we retrospectively analysed the




© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Xin Ni; [nixin@bch.com.cn](mailto:nixin@bch.com.cn)



**Figure 1** The flow diagram of the database search. ICD-10, the 10th revision of the International Classification of Diseases.

epidemiological characteristics of paediatric burn in a large-scale and multicentre study.

Herein, we collected the medical records of 6741 paediatric burn cases from 23 Chinese provincial and municipal medical institutions in the FRCPD database. Excerpt epidemiological data from the medical records were retrieved and analysed. This large-scale, multicentre study can describe the epidemiological characteristics of paediatric burn in China, which can increase child protection against burn injuries, improve the quality of care and reduce hospitalisation costs.

## METHODS

### Data collection and inclusion criteria

In our study, 4350470 medical records were retrospectively investigated from the FRCPD database, and 6741 paediatric burn cases were extracted from 1 January 2016 to 31 December 2019 (figure 1). The following patients were included: (a) the 10th revision of the International Classification of Diseases (ICD-10 code): 'T20-T32 (no T26, T75', 'X00-X19 (no X06, X07, X14)' and 'W85-87, W94, W99'; (b) Only the first hospitalisation of burn children was included. The exclusion criteria were as follow: (a) patients aged >18 years who belonged to general hospitals with adult departments and (b) data of children who had unclear demographic characteristics. The epidemiological characteristics of patients, including gender, age, cause of burn injuries, complication, the timing of hospitalisation (season and month) and hospitalisation length and cost, were collected.

Based on their age, the patients were divided into four categories: <1 year old (infants), 1–2 years old (toddlers), 3–5 years old (preschoolers) and 6–15 years old (school-children). In addition, the cause of burns was grouped into six categories based on the ICD-10 codes: hydrothermal scald (X10-19), flame (W34, W36, W38-40, W92, Y25, Y27, X00, X02, X04-06, X08, X09, X97), electric (W85-87, X33), chemical (X29, X44, X46, X47, X49, X69, Y19, Y41, Y43, Y45, Y46, Y50, Y54, Y55, Y57, Y63),

others (except for above four causes) and unknown (not described).

Fifteen common complications in paediatric burn patients were also collected, such as pneumonia, shock and electrolyte disturbance (the detailed classification is shown in online supplemental materials). According to the hospitalisation season, we divided hospitalisation timing into spring (March–May), summer (June–August), autumn (September–November) and winter (December–February) groups.

### Patient and public involvement

Patients or the public were not involved in the design, or conduct, or reporting, or dissemination plans of our research.

### Statistical analysis

Categorical variables, including gender, age, cause of burn injuries, complication and the timing of hospitalisation, were expressed as numbers (N) and percentages (%) and were compared among different groups using Pearson  $\chi^2$  tests or when there were cells whose count was less than 5 and did not satisfy the requirements of Pearson  $\chi^2$  tests, Fisher's exact tests. Continuous variables included in this study, namely hospitalisation length and cost, were both non-normally distributed according to Shapiro-Wilk tests. Therefore, they were expressed as median and IQRs and compared using Kruskal-Wallis tests. Post-hoc tests were conducted using the Steel-Dwass method. P value < 0.05 was considered statistically significant. Statistical analyses were performed using JMP Pro V.15 software.

## RESULTS

### Gender and age of patients

A total of 6741 medical records were collected. Boys accounted for 63.23% and girls accounted for 36.77% (table 1), with a male-to-female ratio of 1.72:1. Our results showed that boys were markedly more than girls in all age groups and cause of burn injuries groups (figure 2A,B

**Table 1** The baseline characteristics of included paediatric burn patients


Characteristics	Paediatric inpatients (N)	Per cent
<b>Gender</b>		
Male	4262	63.23%
Female	2479	36.77%
<b>Age</b>		
<1 year old	366	5.43%
1–2 years old	4715	69.95%
3–5 years old	1158	17.18%
6–15 years old	502	7.45%
<b>Cause of burn injuries</b>		
Hydrothermal scald	5431	80.57%
Flame	206	3.06%
Electric	85	1.26%
Chemical	84	1.25%
Others	100	1.48%
Unknown	835	12.39%
<b>Surgical intervention</b>		
Yes	1090	16.17%
No	5651	83.83%

and online supplemental table 1,  $p=0.048$  or  $p=0.013$ , respectively).


The age of patients ranged from 1 day to 15 years. As shown in table 1, patients aged 1–2 years were dominant (69.95%), while <1-year-old patients had the prevalence (5.43%).

### Cause of paediatric burn

Among the 6741 hospitalised cases (table 1), the most common cause of burn injuries was hydrothermal scald (80.57%), followed by flame (3.06%), electricity (1.26%), chemicals (1.25%), others (1.48%) and



**Figure 2** The distributions of the hospitalised cases of burn injuries based on the age and gender (A), gender and causes (B) and age and causes (C) of the inpatients, respectively.



**Figure 3** The proportion of common complications in paediatric burn patients.

unknown (12.39%). As shown in online supplemental table 2, the percentage of burn caused by hydrothermal scald was over 50% in different age groups, especially in children aged 1–2 years (85.24%). Thus, the cause distribution was significantly differed with age of the patients ( $p<0.001$ , online supplemental table 2). Moreover, we have drawn the mosaic plot for the age distribution of paediatric burn patients with different causes of burn injuries (figure 2C).

### The complications of paediatric burn

The complications of burn injuries had a great impact on disease prognosis, and many patients suffered from more than one complication. The 15 most common complications were summarised in figure 3. Pneumonia (21%), shock (17%) and electrolyte disturbance (13%) were the top 3 complications of paediatric burn. In addition, table 2 reveals that the age of patients was significantly different with electrolyte disturbance ( $p=0.003$ ), cardiovascular complications ( $p=0.017$ ), hypoproteinaemia ( $p=0.014$ ), intestinal infection ( $p=0.002$ ), aspiration injury ( $p<0.001$ ), hepatic complications ( $p=0.047$ ) and renal complications ( $p<0.001$ ).


### The timing, length and cost of paediatric burn in different age, cause of burn injuries and surgical intervention groups

Regarding the hospitalisation season, our study found that children were mostly admitted to the hospital during spring (26.73%), summer (25.69%), autumn (22.10%) and winter (25.47%), respectively. In particular, the number of paediatric burn patients was minimum in November (7.24%) and maximum in May (9.24%, figure 4 and online supplemental table 3). In the present study, the hospitalisation season distribution was significantly different among the groups of different cause of burn injuries ( $p<0.001$ , online supplemental table 1). Furthermore, the incidence of burn injuries caused by hydrothermal scald was maximum in spring (26.90%), while the flame was more in summer (39.32%), the electric burn was in winter (44.71%) and chemical burn in autumn (32.14%).

**Table 2** The complication and comorbidity distribution of paediatric burn patients in different age groups

Complication and comorbidity	N	Yes or no	<1 year old (n, row per cent)	1–2 years old (n, row per cent)	3–5 years old (n, row per cent)	6–15 years old (n, row per cent)	$\chi^2$	P value
Pneumonia	222	1	16, 7.21%	151, 68.02%	35, 15.76%	20, 9.01%	2.477	0.480
	6519	0	350, 5.37%	4564, 70.01%	1123, 17.23%	482, 7.39%		
Shock	176	1	12, 6.82%	125, 71.02%	29, 16.48%	10, 5.68%	1.480	0.687
	6565	0	354, 5.39%	4590, 69.92%	1129, 17.20%	492, 7.49%		
Electrolyte disturbance	134	1	15, 11.19%	83, 61.94%	20, 14.93%	16, 11.94%	13.731	<b>0.003</b>
	6607	0	351, 5.31%	4632, 70.11%	1138, 17.22%	486, 7.36%		
Septicopyaemia	122	1	13, 10.66%	79, 64.75%	22, 18.03%	8, 6.56%	6.914	0.075
	6619	0	353, 5.34%	4636, 70.04%	1136, 17.16%	494, 7.46%		
Cardiac diseases	117	1	7, 5.98%	70, 59.83%	23, 19.66%	17, 14.53%	10.257	<b>0.017</b>
	6624	0	359, 5.42%	4645, 70.12%	1135, 17.14%	485, 7.32%		
Hypoproteinaemia	82	1	10, 12.20%	49, 59.75%	19, 23.17%	4, 4.88%	10.701	<b>0.014</b>
	6659	0	356, 5.35%	4666, 70.07%	1139, 17.10%	498, 7.48%		
Intestinal infection	68	1	9, 13.24%	53, 77.94%	4, 5.88%	2, 2.94%	15.312	<b>0.002</b>
	6673	0	357, 5.35%	4662, 69.87%	1154, 17.29%	500, 7.49%		
Convulsion	37	1	0, 0.00%	30, 81.08%	5, 13.51%	2, 5.41%	Fisher	0.485
	6704	0	366, 5.46%	4685, 69.88%	1153, 17.20%	500, 7.46%		
Aspiration injury	27	1	3, 11.11%	9, 33.33%	5, 18.52%	10, 37.04%	Fisher	<b>&lt;0.001</b>
	6714	0	363, 5.41%	4706, 70.09%	1153, 17.17%	492, 7.33%		
Alimentary tract haemorrhage and stress ulcer	24	1	1, 4.17%	13, 54.17%	5, 20.83%	5, 20.83%	Fisher	0.083
	6717	0	365, 5.43%	4702, 70.00%	1153, 17.17%	497, 7.40%		
Liver diseases	18	1	4, 22.22%	10, 55.55%	3, 16.67%	1, 5.56%	Fisher	<b>0.047</b>
	6723	0	362, 5.39%	4705, 69.98%	1155, 17.18%	501, 7.45%		
Renal diseases	10	1	1, 10.00%	2, 20.00%	3, 30.00%	4, 40.00%	Fisher	<b>&lt;0.001</b>
	6731	0	365, 5.42%	4713, 70.02%	1155, 17.16%	498, 7.40%		
Multiple organ dysfunction	9	1	0, 0.00%	6, 66.67%	3, 33.33%	0, 0.00%	Fisher	0.629
	6732	0	366, 5.44%	4709, 69.95%	1155, 17.15%	502, 7.46%		
Successful resuscitation from Cardiac arrest	9	1	0, 0.00%	7, 77.78%	2, 22.22%	0, 0.00%	Fisher	0.912
	6732	0	366, 5.44%	4708, 69.93%	1156, 17.17%	502, 7.46%		
Systemic inflammatory response syndrome	8	1	1, 12.50%	7, 87.50%	0, 0.00%	0, 0.00%	Fisher	0.346
	6733	0	365, 5.42%	4708, 69.92%	1158, 17.20%	502, 7.46%		

Bold p values in the table indicate significant differences.

**Figure 4** The hospitalisation month distribution of paediatric burn patients in different age groups.

The median hospitalisation length was 8 days for paediatric burn in this study. There were no significant differences in age groups ( $p=0.065$ , table 3). Interestingly, our results showed that the hospitalisation days significantly differed based on the cause of burn injuries ( $p<0.001$ , table 3). The length of hospitalisation was the longest in patients burned by flame (the average was 10 days). In addition, surgical intervention significantly increased the length of hospital stay, and the median hospitalisation days was 15 days among those who underwent surgery ( $p<0.001$ , table 3).



**Table 3** The hospitalisation days and expenses of paediatric burn patients in different age, cause of burn injuries and surgical intervention groups

	N	Hospitalisation days	$\chi^2$	P value	Hospitalisation expense	$\chi^2$	P value
Age			7.241	0.065		13.807	<b>0.003</b>
<1 year old	366	9 (5, 16)			8364.905 (4726.540, 15 865.008)		
1–2 years old	4715	8 (5, 13)			7651.290 (4340.300, 13 849.540)		
3–5 years old	1158	9 (6, 14)			8197.555 (4563.660, 15 960.208)		
6–15 years old	502	8 (5, 15)			7604.705 (4308.363, 16 412.910)		
Cause of burn injuries			84.825	<0.001		101.697	<b>&lt;0.001</b>
Hydrothermal scald	5431	9 (6, 14)			8087.280 (4544.11, 14 745.15)		
Flame	206	10 (6, 22)			11 333.825 (5341.2275, 32 354.395)		
Electric	85	7 (3, 19)			8295.440 (4180.000, 20 183.535)		
Chemical	84	6 (3, 15)			7360.185 (2867.095, 23 227.935)		
Others	100	8 (4, 12)			6683.780 (3974.323, 11 361.275)		
Unknown	835	7 (5, 11)			5905.230 (3829.800, 10 576.620)		
Surgical intervention			552.466	<0.001		1375.611	<b>&lt;0.001</b>
Yes	1090	15 (9, 26)			24 529.945 (12 745.078, 42 988.745)		
No	5651	8 (5, 12)			6647.910 (4026.300, 11 183.100)		

Bold p values in the table indicate significant differences.

The hospitalisation costs were considered the total expenses of inpatients, including treatment, medications, materials, lab tests, nursing care and additional expenses. **Table 3** revealed that the hospitalisation costs were significantly differed with age ( $p=0.003$ ), cause of burn injuries ( $p<0.001$ ) and surgical intervention ( $p<0.001$ ) among all 6471 cases. The hospitalisation costs were highest in paediatric inpatients with <1 year old or burns caused by flame.

## DISCUSSION

In the current study, we found that among paediatric burn patients, male gender, 1–2 years of age, and hydrothermal scald were more common. Pneumonia was the most common complication in paediatric burn, and the age of patients was significantly differed with burn complications, including electrolyte disturbance, cardiac complications, hypoproteinaemia, intestinal infection, aspiration injury, hepatic complications and renal complications. The number of paediatric burn

patients increased in spring, and the median hospitalisation length was 8 days. Furthermore, the hospitalisation length significantly differed according to surgical intervention and cause of burn injuries. The hospitalisation costs were significant difference with age, surgical intervention and cause of burn injuries.

## Gender and age

Gender and age are significant risk factors for burn injuries. In most developing countries, burn injuries are more common among adult females due to cooking and heating fuels. In contrast, in developed countries, burn injuries are more common among adult males due to industrial accidents.<sup>1</sup> Previous reports revealed that boys are more susceptible to burn injuries because of impulsivity, curiosity, hyperactivity and lack of self-awareness.<sup>10–12</sup> Consistent with these reports, the incidence of burn injuries was much higher in males than females in this study. Furthermore, the number of males who experienced burn injury was much higher than that of females in different age and cause of burn groups.



Paediatric burn primarily occurred in children younger than 5 years old,<sup>13</sup> and our data also showed that children aged 1–2 years were more susceptible. Moreover, accidental burns were less common in infants (<1 year old) who did not walk, preschoolers (3–5 years old) and school children (6–15 years old) who recognised the danger. Given that the age distribution of burned children is wide ranging, education programmes for burn prevention targeting families, schools and child-care facilities are an important public health initiative to reduce the burn burden. For example, it is essential to remind and educate caretakers especially toddler (1–2 years old) caretakers, about burn prevention.<sup>14</sup> Notably, we cannot assume that children aged 15–18 years do not suffer from burn injuries, because children over 14 years are often admitted to adult hospitals in China.

### Cause of burn injuries

Children are susceptible to burn injuries, especially to hydrothermal scald. Various types of burn are related to eating or bathing habits, clothes and environmental temperature. Toddlers are often burned during bathing and eating. For instance, they are usually burned by boiling water, hot soup and hot milk,<sup>10</sup> whereas burn injuries in young adults are often related to flame or electrical injuries.<sup>15–16</sup> In our study, children aged 1–2 years are mainly burned by hydrothermal scald, and school-children aged 6–15 years are more likely burned by flames. Therefore, the necessary preventive measures, such as keeping hot water bottles out of sight of children, or testing the water temperature before bathing, could protect children from burns caused by hydrothermal scalds. In addition, the mental health of adolescents needs serious attention, although only three boys aged 6–15 years with burn injuries caused by intentional self-injury were identified among our cases.

### Complications

In addition, more attention should be paid to complications during hospitalisation. Burn-related complications, such as abdominal compartment syndrome, acute respiratory distress syndrome, septic shock, acute kidney injury and pneumonia, were the leading causes of mortality in paediatric burn.<sup>17</sup> Compared with burns in adults, burns in children aged less than 4 years were easily complicated due to weak skin barriers.<sup>18</sup> Furthermore, severe burn injuries led to organ damage/failure in children.<sup>19</sup> Among the observation in our study, the most common complications was pneumonia. Heat damage and irritants lead to the release of inflammatory mediators in the upper respiratory tract and increased vascular permeability leading to oedema.<sup>20</sup> Currently, the treatment of respiratory injury included antibiotics, airway control and mechanical ventilation. In addition, loss of circulating blood results in hypovolaemic shock, peripheral oedema, multiorgan failure and even death.<sup>21</sup> Children are more susceptible to burn shock than adults due to lower circulating blood volume. According to

the epidemiological information of paediatric burns, medical centres can devote intensive resources to prepare each patient and provide the best service. More effective evidence-based, individualised programmes should be developed to treat paediatric burn with high complication rates.

### Hospitalisation timing, length and cost

The seasonal variation, rather than the yearly variation, seems more statistically valuable in burn injuries. Summer was the peak of severe burn patients admitted to the eight public hospitals.<sup>18</sup> Nevertheless, our study verified that the number of paediatric burn patients was higher in the spring. Most likely, the cold weather limits children's outdoor activities. Hydrothermal solutions and electric sockets are more accessible during indoor activities of children. Inversely, hydrothermal scald injuries are milder when children wear thick clothing, which may partly explain why November had the lowest number of paediatric burn hospitalisation.

Besides, a previous study showed that hospitalisation length is related to the extent of body surface area.<sup>22</sup> A study found that the median hospitalisation length of burn patients increased to 14 days due to severe injury and prolonged rehabilitation.<sup>18</sup> The median hospitalisation length was 8 days in this study. Furthermore, the hospitalisation length was significantly different among groups with different cause of burn and surgical intervention. Paediatric burn caused by flame or surgery may increase median hospitalisation length. Hospitalisation costs were significantly associated with age, surgical intervention and cause of burn injuries. A study found that medications, particularly antibiotics, were the leading cause of hospitalisation costs in paediatric burn.<sup>4</sup> Similarly, treatment and medications were the main costs in this study, and hospitalisation expenses significantly increased in patients <1 year old or undergoing surgery. Our results also confirmed that hospitalisation length and costs of paediatric burn were higher when burn was caused by flame. Therefore, it is essential to reduce the incidence of flame injuries, considering the hospitalisation length and expenses of paediatric burn. Active intervention for flame burns, shortening the interval between two surgeries and increasing the treatment area of a single surgery may reduce the length of hospital stay.

Severity of burns is very important criteria in all type of burn injuries, ultimately affecting morbidity, mortality, complications, hospital stay, hospital expenses and surgical interventions etc. Unfortunately, severity of burns (% total body surface area and depth of burn injuries) in all participants were not analysed in the present study due to incomplete diagnosis of included homepage information. In addition, some children with mild burn injuries received only outpatient treatment, and this might exclude the number of mild burn cases, and influence the distribution of children patients.

## CONCLUSION

Burn injuries are devastating in children, often leading to lifelong physical and psychological impairments. According to the FRCPD database, this large-scale and multi-centre study from 2016 to 2019 revealed the epidemiological characteristics of paediatric burn. Overactive boys aged 1–2 years who lack self-awareness are more likely to suffer from burn, and parents should pay more attention to preventing hydrothermal scald. Pneumonia was the most common complication of burn injuries, which needs more attention and early prevention. Herein, we recommended some measures to protect children from burn injuries, improve treatment and reduce hospitalisation costs in paediatric burn.

### Author affiliations

<sup>1</sup>Department of Burn & Plastic Surgery, Beijing Children's Hospital, Capital Medical University, National Center for Children's Health, Beijing, People's Republic of China

<sup>2</sup>Big Data Center, Beijing Children's Hospital, Capital Medical University, National Center for Children's Health, Beijing, People's Republic of China

<sup>3</sup>Beijing Key Laboratory for Pediatric Diseases of Otolaryngology, Head and Neck Surgery, MOE Key Laboratory of Major Diseases in Children, Beijing Pediatric Research Institute, Beijing Children's Hospital, Capital Medical University, National Center for Children's Health, Beijing, People's Republic of China

<sup>4</sup>Medical Record Management Office, Beijing Children's Hospital, Capital Medical University, National Center for Children's Health, Beijing, People's Republic of China

<sup>5</sup>Information Center, Beijing Children's Hospital, Capital Medical University, National Center for Children's Health, Beijing, People's Republic of China

<sup>6</sup>Department of Otolaryngology Head and Neck Surgery, Beijing Children's Hospital, Capital Medical University, National Center for Children's Health, Beijing, People's Republic of China

**Contributors** The authors' contributions were as follows. YG and XN contributed to the conception and design of the study. KC, JL, XW contributed to acquisition, analysis and interpretation of the data. KC and JL wrote the manuscript. PC, GF, YZ, XX revised the manuscript. XN acted as a guarantor in the manuscript. All authors read and approved the final manuscript.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not applicable.

**Ethics approval** All patient records were pseudonymised by the data providers through encryption of personal identifiers. Ethical approval to hold and analyse the data set was obtained from the ethics committee of Beijing Children's Hospital, Capital Medical University (Approval Number: 2020-k-10).

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

### ORCID iD

Xin Ni <http://orcid.org/0000-0002-7781-2600>

## REFERENCES

- 1 Puthumana JS, Ngaage LM, Borrelli MR, *et al*. Risk factors for cooking-related burn injuries in children, who global burn registry. *Bull World Health Organ* 2021;99:439–45.
- 2 Atiyeh BS, Costagliola M, Hayek SN. Burn prevention mechanisms and outcomes: pitfalls, failures and successes. *Burns* 2009;35:181–93.
- 3 Mukerji G, Chamanian S, Patidar GP, *et al*. Epidemiology of paediatric burns in indore, India. *Burns* 2001;27:33–8.
- 4 Burd A, Yuen C. A global study of hospitalized paediatric burn patients. *Burns* 2005;31:432–8.
- 5 Zhu L, Zhang Y, Liu L, *et al*. Hospitalized pediatric burns in North China: a 10-year epidemiologic review. *Burns* 2013;39:1004–11.
- 6 Hai Jun W, Jie X, Jun Z, *et al*. Comparable results of epidemiology of children with burns among different decades in a burn unit in Jinzhou, China. *Burns* 2011;37:513–20.
- 7 Xin W, Yin Z, Qin Z, *et al*. Characteristics of 1494 pediatric burn patients in Shanghai. *Burns* 2006;32:613–8.
- 8 Wang X, Zeng Y, Tian J, *et al*. A brief introduction to the futang updating medical records (future) database. *Pediatr Investig* 2021;5:247–8.
- 9 Feng G, Zeng Y, Tian J, *et al*. Disease spectrum analysis of hospitalized children in China: a study of 18 tertiary children's hospitals. *Pediatr Investig* 2019;3:159–64.
- 10 Dissanaik S, Boshart K, Coleman A, *et al*. Cooking-related pediatric burns: risk factors and the role of differential cooling rates among commonly implicated substances. *J Burn Care Res* 2009;30:593–8.
- 11 Drago DA. Kitchen scalds and thermal burns in children five years and younger. *Pediatrics* 2005;115:10–6.
- 12 Rayner R, Prentice J. Paediatric burns: a brief global review. *Wound Pract Res* 2011;11:39–46.
- 13 Dissanaik S, Rahimi M. Epidemiology of burn injuries: highlighting cultural and socio-demographic aspects. *Int Rev Psychiatry* 2009;21:505–11.
- 14 Parbhoo A, Louw QA, Grimmer-Somers K. Burn prevention programs for children in developing countries require urgent attention: a targeted literature review. *Burns* 2010;36:164–75.
- 15 Taylor AJ, McGwin G, Cross JM, *et al*. Serious occupational burn injuries treated at a regional burn center. *J Burn Care Rehabil* 2002;23:244–8.
- 16 Maghsoudi H, Gabraely N. Epidemiology and outcome of 121 cases of chemical burn in East azarbaijan Province, Iran. *Injury* 2008;39:1042–6.
- 17 Aldekhayel S, Khubrani AM, Alshaalan KS, *et al*. Outcomes and complications of diabetic burn injuries: a single center experience. *Int J Burns Trauma* 2021;11:220–5.
- 18 Tian H, Wang L, Xie W, *et al*. Epidemiologic and clinical characteristics of severe burn patients: results of a retrospective multicenter study in China, 2011–2015. *Burns Trauma* 2018;6:14.
- 19 Porter C, Herndon DN, Sidossis LS, *et al*. The impact of severe burns on skeletal muscle mitochondrial function. *Burns* 2013;39:1039–47.
- 20 Nielson CB, Duethman NC, Howard JM, *et al*. Burns: pathophysiology of systemic complications and current management. *J Burn Care Res* 2017;38:e469–81.
- 21 Rae L, Fidler P, Gibran N. The physiologic basis of burn shock and the need for aggressive fluid resuscitation. *Crit Care Clin* 2016;32:491–505.
- 22 Karimi H, Montealvan A, Motabar AR, *et al*. Epidemiology of paediatric burns in Iran. *Ann Burns Fire Disasters* 2012;25:115–20.

**The detailed classification of the complications in pediatric burn patients:**

- (1) Pneumonia: whooping cough, lung infection, mycoplasma pneumonia, community-acquired pneumonia, severe pneumonia.
- (2) Shock: hypovolemic shock, hemorrhagic shock, traumatic shock, septic shock, lightning shock, toxic shock.
- (3) Electrolyte disturbance: low sodium, high sodium, low potassium, high potassium, low chloride, high chlorine, low calcium, high calcium, metabolic acidosis, dehydration.
- (4) Septicopyemia: sepsis, bacteremia.
- (5) Cardiac diseases: myocardial damage, myocardial strain, heart failure.
- (6) Hypoproteinemia: albumin abnormality.
- (7) Intestinal infection: Diarrhea, enteritis, bacterial intestinal infection, rotavirus enteritis.
- (8) Convulsion: febrile convulsions, seizure.
- (9) Aspiration injury.
- (10) Alimentary tract hemorrhage; stress ulcer.
- (11) Liver diseases: abnormal liver function, liver insufficiency, liver injury, liver failure, hepatic coma.
- (12) Renal diseases: Renal insufficiency, renal failure, tubular necrosis.
- (13) Multiple organ dysfunction: multiple organ failure; multiple organ dysfunction syndrome.
- (14) Successful resuscitation from cardiac arrest.
- (15) Systemic inflammatory response syndrome.



**Sup Table 1** Gender distribution of pediatric burn patients in different age and cause of burn injuries groups.

	N	Male (n, row percent)	Female (n, row percent)	$\chi^2$	P
<b>Age</b>				7.905	<b>0.048</b>
<1 years old	366	211, 57.65%	155, 42.35%		
1-2 years old	4715	2992, 63.46%	1723, 36.54%		
3-5 years old	1158	724, 62.52%	434, 37.48%		
6-15 years old	502	335, 66.73%	167, 33.27%		
<b>Cause of burn injuries</b>				14.544	<b>0.013</b>
Hydrothermal scald	5431	3399, 62.59%	2032, 37.41%		
Flame	206	137, 66.50%	69, 33.50%		
Electric	85	69, 81.18%	16, 18.82%		
Chemical	84	53, 63.10%	31, 36.90%		
Others	100	66, 66.00%	34, 34.00%		
Unknown	835	538, 64.43%	297, 35.57%		

**Sup Table 2** Cause distribution of pediatric burn patients in different age groups.

	N	<1 years old (n, column percent)	1-2 years old (n, column percent)	3-5 years old (n, column percent)	6-15 years old (n, column percent)	$\chi^2$	P
<b>Cause of burn injuries</b>						665.049	<b>&lt;0.001</b>
Hydrothermal scald	5431	254, 69.40%	4019, 85.24%	882, 76.17%	276, 54.98%		
Flame	206	8, 2.19%	62, 1.31%	46, 3.97%	90, 17.93%		
Electric	85	1, 0.27%	31, 0.66%	28, 2.42%	25, 4.98%		
Chemical	84	3, 0.82%	49, 1.04%	17, 1.47%	15, 2.99%		
Others	100	8, 2.19%	55, 1.17%	18, 1.55%	19, 3.78%		
Unknown	835	92, 25.14%	499, 10.58%	167, 14.42%	77, 15.34%		

**Sup Table 3** Age distribution of pediatric burn patients in different hospitalization

month.

Month	Total (N, percent)	<1 years old	1-2 years old	3-5 years old	6-15 years old
January	596, 8.84%	41	380	129	46
February	522, 7.74%	15	315	128	64
March	588, 8.72%	7	405	131	45
April	589, 8.74%	8	441	108	32
May	623, 9.24%	13	458	111	41
June	602, 8.93%	11	458	94	39
July	587, 8.71%	15	431	91	50
August	560, 8.31%	26	403	79	52
September	541, 8.03%	26	408	75	32
October	505, 7.49%	48	356	69	32
November	488, 7.24%	71	320	68	29
December	540, 8.01%	85	340	75	40

**Sup Table 4** Hospitalization season distribution of pediatric burn patients in different cause of burn injuries groups.

	N	Hydrother mal scald (n, row percent)	Flame (n, row percent)	Electric (n, row percent)	Chemica l (n, row percent)	Others (n, row percent)	Unknow n (n, row percent)	$\chi^2$	P
<b>Hospital ization season</b>								68.610	<0.001
Spring	1802	1461, 26.90%	47, 22.82%	18, 21.28%	21, 25.00%	22, 22.00%	233, 27.90%		
Summer	1732	1333, 24.54%	81, 39.32%	7, 8.24%	26, 30.95%	30, 30.00%	240, 28.74%		
Autumn	1490	1205, 22.19%	35, 16.99%	22, 25.88%	27, 32.14%	23, 23.00%	178, 21.32%		
Winter	1717	1432, 26.37%	43, 20.87%	38, 44.71%	10, 11.90%	25, 25.00%	184, 22.04%		