



# Vitamin D insufficiency in infants with increased risk of developing type 1 diabetes: a secondary analysis of the POInT Study

An Jacobs <sup>1</sup>, Maarten Warnants,<sup>2</sup> Veronika Vollmuth,<sup>3</sup> Christiane Winkler,<sup>3,4</sup> Andreas Weiss,<sup>3</sup> Anette-Gabriele Ziegler,<sup>3,5</sup> Markus Lundgren,<sup>6,7</sup> Helena Elding Larsson,<sup>8,9</sup> Olga Kordonouri <sup>10</sup>, Thekla von dem Berge,<sup>10</sup> Marie-Luise Zielmann,<sup>11</sup> Ezio Bonifacio,<sup>12</sup> Angela Hommel,<sup>12</sup> Mariusz Oltarzewski,<sup>13</sup> Agnieszka Szypowska,<sup>14</sup> Rachel Besser,<sup>15,16</sup> John A. Todd,<sup>16</sup> Kristina Casteels<sup>1,17</sup>

**To cite:** Jacobs A, Warnants M, Vollmuth V, *et al.* Vitamin D insufficiency in infants with increased risk of developing type 1 diabetes: a secondary analysis of the POInT Study. *BMJ Paediatrics Open* 2024;**8**:e002212. doi:10.1136/bmjpo-2023-002212

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjpo-2023-002212>).

Received 2 August 2023  
Accepted 7 November 2023



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

**Correspondence to**  
Dr An Jacobs; [an.k.jacobs@gmail.com](mailto:an.k.jacobs@gmail.com)

## ABSTRACT

**Background** Vitamin D insufficiency (VDI) may be a factor in the development of type 1 diabetes (T1D). The aim of this study is to investigate the presence and persistence of VDI in a large cohort of infants with increased risk of developing T1D, in light of the differences in local supplementation guidelines.

**Methods** In the POInT Study, a multicentre primary prevention study between February 2018 and March 2021 in Germany, Poland, Belgium, England and Sweden, including infants aged 4–7 months at high genetic risk of developing  $\beta$ -cell autoantibodies, vitamin D levels were analysed at each study visit from inclusion (4–7 months) until 3 years, with an interval of 2 months (first three visits) or 4–6 months (visits 4–8). The protocol actively promotes vitamin D sufficiency to optimise immune tolerance. VDI was defined as a concentration below 30 ng/mL and was treated according to local guidelines of participating centres. Recovery from VDI was defined as a concentration above or equal to 30 ng/mL on the subsequent visit after VDI.

**Results** 1050 infants were included, of which 5937 vitamin D levels were available for analyses. VDI was observed in 1464 (24.7%) visits and 507 (46.1%) of these were not resolved at the next visit. The risk of having VDI was independently associated with season (higher in winter), weight (higher with increased weight), age (higher with increased age) and country (higher in England). The risk of not recovering from VDI was independently associated with the season of the previously determined VDI, which was higher if VDI was identified in winter.

**Conclusions** VDI is frequent in infants with increased risk of developing T1D. Treatment guidelines for VDI do not seem effective. Increasing supplementation dosages in this patient population seems warranted, especially during winter, and increasing dosages more aggressively after VDI should be considered.

## INTRODUCTION

Vitamin D can be obtained from ultraviolet (UV)B-dependent endogenous production, from the diet and from supplements.<sup>1</sup> When

## WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Vitamin D supplementation guidelines for infants and children differ greatly per country.
- ⇒ The association between vitamin D insufficiency and type 1 diabetes is well established, although a causative role in type 1 diabetes development remains to be proven.
- ⇒ In the POInT Study, including children with an increased genetic risk of developing type 1 diabetes, vitamin D levels are systematically analysed, and parents are counselled to adjust supplementation doses accordingly to avoid vitamin D insufficiency.

## WHAT THIS STUDY ADDS

- ⇒ Vitamin D insufficiency is frequent and persistent in infants and children with an increased genetic risk of developing type 1 diabetes, despite systematic counselling.
- ⇒ Even with large increments in supplementation dose after vitamin D insufficiency, vitamin D intoxication is extremely rare in the study population.

## HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ Adaptation of the guidelines in terms of standard dose of vitamin D supplementation, as well as dose increments after vitamin D insufficiency, should be considered in this population.
- ⇒ Factors such as season and body mass index for age could be taken into account when treating vitamin D insufficiency in this group of children with increased type 1 diabetes risk.

UVB hits the skin, it leads to the production of pre-vitamin D, which is an unstable intermediate thermally converted into vitamin D<sub>3</sub>. Vitamin D may also be obtained to a lesser extent from the diet or from vitamin D supplements. Importantly, both synthesised and dietary-obtained vitamin D is an inactive



prohormone and both 25-hydroxylation and 1 $\alpha$ -hydroxylation are required for activation. The 25-hydroxylation occurs in the liver and 25-OH-vitamin D reflects the nutritional vitamin D status. The next step to obtain active vitamin D is 1 $\alpha$ -hydroxylation in the kidneys.<sup>2</sup>

There is controversy about the level of adequacy of vitamin D. The Endocrine Society defines deficiency as 25-(OH)D<sub>3</sub> levels less than 20 ng/mL (50 nmol/L) and vitamin D insufficiency (VDI) as levels 21–29 ng/mL (52–72 nmol/L).<sup>3</sup> The Institute of Medicine defines deficiency as 25-(OH)D<sub>3</sub> levels less than 12 ng/mL (30 nmol/L) and insufficiency as levels 12–20 ng/mL (30–50 nmol/L).<sup>4</sup> A recent French expert consensus paper recommends a 25(OH)D level >20 ng/mL (50 nmol/L) in children to prevent rickets and a 25(OH)D level >30 ng/mL (75 nmol/L) to avoid any mineralisation defects.<sup>5</sup> However, these cut-offs have been defined based on associations with bone metabolism, with markers such as parathyroid hormone and bone fracture risk,<sup>6,7</sup> whereas much higher vitamin D levels are needed to influence the immune effects of vitamin D in animal studies.<sup>2,8</sup> Low levels of vitamin D are common, as a study in 6275 American children and adolescents aged 1–21 years showed that 61% were 25-(OH)D<sub>3</sub> insufficient (15–29 ng/mL) and 9% deficient (<15 ng/mL).<sup>9</sup> When treating VDI, there is a risk of reaching a vitamin D intoxication with concentrations of more than 150 ng/mL (374 nmol/L), which can lead to hypercalcaemia and an imbalance in the regulation of bone metabolism.<sup>1</sup>

There is also controversy and discussion about the correct supplementation regime. The Endocrine Society recommends the following for vitamin D intake: 400–1000 IU, 600–1000 IU and 1500–2000 IU daily for children under 1 year, children 1–18 years and all adults, respectively, to treat and prevent vitamin D deficiency.<sup>10</sup> The expert consensus paper recommends supplementing healthy children 0–18 years of age with a minimum of 400 IU and a maximum of 800 IU of vitamin D per day.<sup>5</sup>

The importance of vitamin D in the regulation of both the innate and adaptive immune system was demonstrated by the discovery of the presence of VDR expression in almost all cells of the immune system.<sup>2,11</sup> Essentially, adequate levels of vitamin D result in a shift in the immune status towards a more tolerogenic status.<sup>12</sup> This makes vitamin D an appealing immune modulator in the treatment of immune-mediated diseases. Epidemiological data showed an association between 25-(OH)D<sub>3</sub> deficiency and the incidence of type 1 diabetes (T1D),<sup>13,14</sup> but that VDI is a causal factor in T1D development remains unproven.<sup>15,16</sup>

Studies in animal models of autoimmune diseases showed that restoring serum 25-(OH)D<sub>3</sub> levels using high doses of vitamin D metabolites or analogues could alter the course of autoimmune diseases like T1D, multiple sclerosis or rheumatoid arthritis.<sup>17</sup> In humans, the European Diabetes Centre's EURODIAB Study showed a significant decrease in T1D diagnosis in children receiving vitamin D supplementation,<sup>18</sup> while other studies indicated that

maternal vitamin D supplementation during pregnancy did not reduce the risk of T1D in the offspring.<sup>19,20</sup> In a recent study of 395 children with T1D between 3 and 18 years old without vitamin D supplementation, 64% were vitamin D insufficient.<sup>21</sup>

The goal of this subanalysis of the Primary Oral Insulin Trial (POInT Study) was to investigate the presence of VDI and the evolution of these insufficiencies over time, and to analyse the effect of country-specific supplementation guidelines and participant characteristics.

## METHODS

The POInT Study, a multicentre primary prevention study that aims to induce immune tolerance to  $\beta$ -cell autoantigens through regular exposure to oral insulin, also aimed for vitamin D sufficiency to optimise the immune response, and therefore, vitamin D levels were repeatedly measured (registered as NCT03364868 on ClinicalTrials.gov). All infants aged 4–7 months with a high genetic risk of developing  $\beta$ -cell autoantibodies were eligible for inclusion and were recruited in the seven participating centres: Munich, Hanover and Dresden (Germany), Warsaw (Poland), Malmö (Sweden), Oxford (England) and Leuven (Belgium). Exclusion criteria were concomitant disease or treatment that may interfere with the assessments, any condition that could be associated with poor compliance or may jeopardise the infant's safe participation, diagnosis of diabetes at the time of recruitment or participation in another clinical trial.<sup>22</sup> For this secondary analysis, visits without a vitamin D measurement were excluded. Patients were not involved in the study design but in the prioritisation of the research question of T1D prevention. Patients support recruitment through dissemination and participation in press conferences. The participating families previously assessed the burden of the intervention in a prior pilot prevention trial.<sup>22,23</sup>

25-OH-vitamin D levels were analysed at the local laboratory of the participating centres at each study visit from the age of 4–7 months until 3 years of age, with an interval of 2 months (for the first three visits) or 4–6 months (from visit 4 to 8). VDI was defined as a vitamin D concentration below 30 ng/mL. Study nurses and physicians at each site were notified via email by the centralised study software if vitamin D values were too low. VDI was then subsequently treated according to the local guidelines of the participating centres, which is shown in [table 1](#). Parents were instructed to continue the same supplementation dose if they were not contacted after the study visit, as they were not actively informed when vitamin D levels were sufficient. Recovery from VDI was defined as a concentration above or equal to 30 ng/mL, the subsequent visit after VDI.

Multivariable logistic regression analysis, adjusted for season (January–March as winter, April–June as spring, July–September as summer, October–December as autumn), sex, history of T1D in a first-degree relative,

**Table 1** Overview of local vitamin D supplementation guidelines per participating centre

	Standard dosage <1 year	Standard dosage >1 year
Germany	500 IU/day	500 IU/day
Belgium	400 IU/day	400 IU/day
England	400 IU/day	400 IU/day
Poland	400–600 IU/day	600–1000 IU/day
Sweden	400 IU/day	400 IU/day
	If vitamin D <30 µg/L <1 year	If vitamin D <30 µg/L >1 year
Germany	Plus 500 IU increase	Plus 1000 IU increase
Belgium	Plus 66 IU increase	Plus 66 IU increase
England	Plus 600 IU increase	Plus 600 IU increase
Poland	Plus 300–600 IU increase	Plus 500–1000 IU increase
Sweden	Plus 400–800 IU increase	Plus 400–2800 IU increase

age, weight, height and country, was used to assess independent associations with the risk of VDI and the chance of recovery from VDI. Effect sizes are reported as ORs and 95% CIs. Statistical analyses were performed with JMP Pro V.14.0.0 (SAS Institute, Cary, North Carolina, USA). Statistical significance was set at a p value of <0.05.

## RESULTS

In total, 1050 children were included, of which 5937 vitamin D levels were available for analyses. Baseline characteristics of these participants are shown in [table 2](#).

[Figure 1](#) represents the distribution of the vitamin D levels per country. The number of patients and the available vitamin D levels per visit per country are illustrated in online supplemental figure 1A,B. Of the 5937 measured vitamin D levels, 1464 (24.7%) were below 30 ng/mL and defined as VDI. The number of children who had at least one VDI at any time point was 667 out of the 1050 (63.5%). Of the 1464 measured VDIs, 592 (53.9%) insufficiencies rose above the threshold of 30 ng/mL at the next visit, whereas 507 insufficiencies (46.1%) were not resolved at the next visit. Of the 592 recovered insufficiencies, 33 (5.6%) were in the excess range above 50 ng/mL, and none were in the toxic range above 150 ng/mL. In this subgroup of the 592 recovered insufficiencies, there was no difference between the participating countries in normal or excess ranges ( $p=0.66$ ). 4476 out of the total of 5937 measurements were defined as vitamin D sufficiency (75.3%), and 686 of them (15.3%) were classified as a VDI on the next visit.


### Risk of having a VDI

In multivariable logistic regression, the risk of having VDI was independently associated with the season, weight, age and country ([table 3](#)). The risk of VDI was higher

**Table 2** Baseline characteristics of the participants

	Participants N=1050
Male sex, N (%)	527 (50.3)
Age (months) at visit 1, median (IQR)	5.9 (5.2–6.5)
Weight (kg) at visit 1, median (IQR)	7.8 (7.1–8.4)
Height (cm) at visit 1, median (IQR)	68 (66–69)
Positive history of T1DM in first-degree relatives, N (%)	556 (53.0)
Vitamin D level (ng/mL) of all visits combined, median (IQR)	36 (30–43)
Vitamin D level (ng/mL) of all visits combined, mean (SD)	37 (11)
Vitamin D toxic levels (>150 ng/mL) of all visits combined, N (%)	1 (0.02)
Country, N (%)	
Germany	504 (48.0)
Belgium	80 (7.6)
England	51 (4.9)
Poland	242 (23.0)
Sweden	173 (16.5)
Participants with vitamin D level available on each visit, N (%)	
Visit 1	965 (91.9)
Visit 2	957 (91.1)
Visit 3	937 (89.2)
Visit 4	930 (88.6)
Visit 5	811 (77.2)
Visit 6	641 (61.0)
Visit 7	443 (42.2)
Visit 8	253 (24.1)
T1DM, type 1 diabetes mellitus.	

in winter as compared with all other seasons, and lower in summer as compared with all other seasons. Also, for every kilogram increase in body weight and for every year increase in age, the risk of VDI increased. Finally, the risk of having VDI was higher in England as compared with Germany (OR 1.42 (1.06 to 1.90),  $p=0.01$ ), Poland (OR 1.69 (1.24 to 2.31),  $p=0.0009$ ) and Belgium (OR 1.43 (1.00 to 2.03),  $p=0.04$ ). The risk was lower in Poland as compared with Germany (OR 0.83 (0.71 to 0.99),  $p=0.03$ ) and Sweden (OR 0.79 (0.65 to 0.99),  $p=0.03$ ). There was no independent association between height, sex or a first-degree relative with T1D and the risk of VDI. The repeated measurements analysis ([table 4](#)), which corrects for multiple measurements of vitamin D of the same child, shows similar results for season, age and body weight. However, the effect of country largely disappears, which indicates that this effect is probably mostly driven by children who have persistent insufficiencies.



**Figure 1** Violin plots of the vitamin D levels per country. The kernel density estimation visualises the distribution of the data. Each blue dot represents a single vitamin D level.

### Risk of not recovering from a VDI

The risk of not recovering from VDI was independently associated with the season of the previously determined VDI (table 5). If the VDI was identified in winter, the risk of not recovering was higher as compared with insufficiencies identified in spring or summer. Insufficiencies identified in summertime had higher chances of recovery than the insufficiencies in autumn or spring. None of the other available variables was associated with the risk of not recovering from VDI.

### DISCUSSION

This study covers a large sample size of almost 6000 analysed 25-OH-vitamin D levels in 1050 infants and children between the age of 4 months and 3 years with an increased genetic risk of developing  $\beta$ -cell autoantigens, in five European countries. One-fourth of the measured values were defined as at least a VDI, and almost half of these insufficiencies were not resolved by the next visit. The risk of having a VDI increased in winter, with higher age and body weight, and in England. A risk factor for not recovering from a VDI was if the VDI was measured in wintertime.

Since VDI is associated with T1D and might play a role in the pathogenesis,<sup>24</sup> and since VDI is associated with rickets, muscle weakness and other chronic diseases,<sup>25</sup> its prevalence and its persistence after dose increments in this population of infants with a confirmed increased genetic risk of developing T1D are unexpected. However, in a substudy of the TRIGR Study, investigating the effect of an extensively hydrolysed formula on the incidence of T1D,<sup>16</sup> the mean vitamin D levels at baseline in infants (until 18 months) with an increased genetic risk of T1D with and without autoantibodies were 26.2 ng/mL and 26.9 ng/mL, respectively.<sup>26</sup> The mean vitamin D levels during follow-up in our study are significantly higher (37 ng/mL), which might be explained by the regularity of the blood analyses, the close follow-up and emphasis on the importance of compliance towards the parents. However, despite this systemic counselling, the prevalence of VDI was still 25%, which is comparable with the prevalence of VDI in healthy infants: depending on the geographical population that was examined, the prevalence of a VDI is ranging from 19.3% to 40%.<sup>27–29</sup> Moreover, although recommendations on increasing the vitamin D supplementation when necessary were also systematically made,

**Table 3** Multivariable logistic regression analysis of the risk of a vitamin D insufficiency

	OR (95% CI)	P value
Season		<0.0001
Summer vs winter	0.42 (0.35 to 0.50)	<0.0001
Autumn vs winter	0.67 (0.56 to 0.80)	<0.0001
Spring vs winter	0.77 (0.65 to 0.91)	0.002
Weight		
Per kg increase	1.09 (1.03 to 1.16)	0.002
Height		
Per cm increase	0.99 (0.97 to 1.01)	0.56
Age		
Per year increase	1.03 (1.01 to 1.05)	0.003
Country		0.008
England vs Germany	1.42 (1.06 to 1.90)	0.01
Poland vs Germany	0.84 (0.71 to 0.99)	0.03
Belgium vs Germany	1.00 (0.78 to 1.27)	0.96
Sweden vs Germany	1.05 (0.87 to 1.26)	0.60
Gender		
Male vs female	1.03 (0.91 to 1.17)	0.58
First-degree relative (FDR) with T1DM		
FDR vs no FDR	1.03 (0.90 to 1.17)	0.60
T1DM, type 1 diabetes mellitus.		

**Table 4** Repeated measurements analysis of the risk of a vitamin D insufficiency

	OR (95% CI)	P value
Season		<0.0001
Summer vs winter	0.38 (0.31 to 0.47)	<0.0001
Autumn vs winter	0.64 (0.52 to 0.78)	<0.0001
Spring vs winter	0.54 (0.45 to 0.65)	0.006
Weight		
Per kg increase	1.11 (1.03 to 1.20)	0.008
Height		
Per cm increase	0.99 (0.97 to 1.02)	0.61
Age		
Per year increase	1.03 (1.01 to 1.06)	0.009
Country		0.06
England vs Germany	1.53 (1.00 to 2.34)	0.05
Poland vs Germany	0.80 (0.63 to 1.01)	0.06
Belgium vs Germany	1.00 (0.71 to 1.43)	0.99
Sweden vs Germany	1.04 (0.79 to 1.35)	0.79
Gender		
Male vs female	1.04 (0.87 to 1.26)	0.64
First-degree relative (FDR) with T1DM		
FDR vs no FDR	1.03 (0.86 to 1.25)	0.71
T1DM, type 1 diabetes mellitus.		

**Table 5** Multivariable logistic regression analysis of the risk of not recovering from a vitamin D insufficiency (VDI)

	OR (95% CI)	P value
Season of the first VDI measured		<0.0001
Summer vs winter	0.37 (0.26 to 0.54)	<0.0001
Autumn vs winter	0.67 (0.44 to 1.03)	0.06
Spring vs winter	0.65 (0.46 to 0.93)	0.01
Weight		
Per kg increase	1.04 (0.91 to 1.18)	0.52
Height		
Per cm increase	1.00 (0.95 to 1.05)	0.85
Age		
Per year increase	0.99 (0.95 to 1.03)	0.66
Country		0.65
Gender		
Male vs female	1.03 (0.79 to 1.33)	0.81
First-degree relative (FDR) with T1DM		
FDR vs no FDR	0.96 (0.74 to 1.25)	0.79
T1DM, type 1 diabetes mellitus.		

almost half of the insufficiencies were not resolved at the next study visit. Remarkably, having a first-degree family member with T1D, which was the case in more than half of the children, and which could potentially influence adherence, did not affect the risk of a VDI or of a VDI recovery.

Taking into account the results of the multivariable analysis, we can speculate on how to improve the clinical approach in these patients to avoid VDI and to resolve their insufficiency faster. First, as the supplementation guidelines differ between the participating countries, and as a country such as Poland with a higher range and maximum of supplementation guidelines is independently associated with a lower risk of a VDI, it seems warranted to increase the supplementation dosage. Only 1 of nearly 6000 vitamin D levels reached toxic levels (>150 ng/mL). Moreover, although not decreasing the risk of VDI recurrences, the largest range of increasing supplementation dosages after insufficiency in the local guidelines in our study was in Sweden (depending on the severity of insufficiency: 400–800 IU/day for infants younger than 1 year of age, 400–2800 IU/day for older children). Therefore, it seems safe to increase the supplementation dosages in countries where they are low in comparison with Sweden. Next, systematically increasing the dosage in wintertime should be considered in these children. Conversely, more practically challenging is increasing supplementation dosages with increased weight or body mass index for age. A recent randomised controlled trial showed that daily 6000 IU vitamin D supplementation in obese children was safe and sufficient to reach vitamin D sufficiency.<sup>30</sup> Increasing

more aggressively in obese children could be taken into account when increasing dosages in these children does not resolve a previously identified vitamin D deficit.

Finally, the strategy of increasing the dosage, which differed extensively between countries, when a VDI was measured, did not contribute to the risk of not resolving the VDI. However, as almost half of the insufficiencies were not resolved after increasing the dose, increasing more aggressively in wintertime seems warranted, as this was the only independent contributor to the risk of not recovering. As shown in a recent randomised controlled trial in children and adolescents with newly diagnosed T1D, even very high dosages of adjunctive ergocalciferol supplementation in high dosage (50 000 IU weekly) was safe and did not lead to toxic levels.<sup>31</sup>

Limitations in this study include the lack of data available on the actual administered dose of vitamin D supplementation by the parents. The use of diaries or other measures of adherence could have improved the analysis. Additionally, quantification of the 25-OH-vitamin D levels was not centralised but was performed by the local laboratories of the participating centres. Since all local laboratories are certified, we believe that the risks of this affecting the results are minimal. The strengths of the study include the multicentre design, as well as the large cohort of more than 1000 infants with increased risk of T1D without missing data for the included variables, of which nearly 6000 vitamin D levels were available for analysis.

## CONCLUSION

Despite the established health benefits of having sufficient levels of vitamin D and our emphasis to parents on the importance of adequate supplementation, VDI is frequent in infants and young children with increased risk of developing T1D, as seen in children not selected on high-genetic T1D risk. The insufficiency remains unresolved very often, even after using treatment dosages. Increasing supplementation dosages in this patient population seems warranted, especially during wintertime. When VDI is detected, increasing dosages more aggressively should be considered.

### Author affiliations

<sup>1</sup>Department of Pediatric Endocrinology and Diabetes, KU Leuven University Hospitals Leuven, Leuven, Belgium

<sup>2</sup>Katholieke Universiteit Leuven, Leuven, Belgium

<sup>3</sup>Institute of Diabetes Research, Helmholtz Munich, German Center for Environmental Health, Munich, Germany

<sup>4</sup>Forschergruppe Diabetes at Klinikum rechts der Isar, School of Medicine, Technical University Munich, Munich, Germany

<sup>5</sup>Forschergruppe Diabetes, Klinikum rechts der Isar, Technische Universität München, Munich, Germany

<sup>6</sup>Department of Clinical Sciences Malmö, Lund University, Malmö, Sweden

<sup>7</sup>Department of Pediatrics, Kristianstad Hospital, Kristianstad, Sweden

<sup>8</sup>Department of Pediatrics, Skåne University Hospital, Malmö, Sweden

<sup>9</sup>Department of Clinical Sciences Malmö, Lund University, Lund, Sweden

<sup>10</sup>Auf der Bult Children's Hospital, Hannover, Germany

<sup>11</sup>Department of Pediatrics, Faculty of Medicine and University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany

<sup>12</sup>Center for Regenerative Therapies, Technische Universität Dresden, Dresden, Germany

<sup>13</sup>Department of Screening and Metabolic Diagnostics, Institute of Mother and Child, Warsaw, Poland

<sup>14</sup>Department of Paediatrics, Medical University of Warsaw, Warsaw, Poland

<sup>15</sup>Department of Paediatrics, University of Oxford, Oxford, UK

<sup>16</sup>Wellcome Centre for Human Genetics, Nuffield Department of Medicine, NIHR Oxford Biomedical Research Centre, University of Oxford, Oxford, UK

<sup>17</sup>Department of Development and Regeneration, KU Leuven, Leuven, Belgium

**Acknowledgements** POInT Study group: GPPAD-Coordinating Centre (CC): Melanie Gündert, Florian Haupt, Stefanie Arnolds, Miriam Bißbort, Karina Blasius, Nadine Friedl, Cigdem Gezgin, Gertrud Göppel, Martin Heigermoser, Maja Jergl, Bianca Höfelschweiger, Manja Jolink, Krisztina Kisfűgedi, Nadine Klein, Ramona Lickert, Claudia Matzke, Kim Muñoz Alvarez, Rebecca Niewöhner, Marlon Scholz, Katharina Schütte-Borkovec, Andreas Weiß, José Maria Zapardiel Gonzalo, Tarini Singh, Sarah Schmidt, Philipp Sift, Merve Vurucu, Katharina Sarcletti, Melanie Sporreiter, Stefanie Jacobson, Ivo Zeller; Medical Monitor: Katharina Warncke; Outcome Committee: Peter Achenbach, Ezio Bonifacio; Type 1 diabetes endpoint committee: Helena Elding Larsson, Anette-G. Ziegler; Pharmacovigilance Committee: Peter Achenbach, Katharina Schütte-Borkovec, Anette-G. Ziegler; Belgium Clinical Centre: Kristina Casteels, Charlien Janssen, Anne Rochtus, Hilde Morobé, Brontë Vrancken, Natalie Van den Driessche, Gert Van Poel, Renka Van Heyste, Janne Houben, Veerle Vanhuysse; Germany, Dresden Clinical Centre: Ezio Bonifacio, Reinhard Berner, Sari Arabi, Lisa Barbknecht, Sevina Dietz, Franziska Ehrlich, Gita Gemulla, Zahra Gholizadeh, Raphael Hoffmann, Angela Hommel, Franziska Lange, Anja Loff, Robert Morgenstern, Anne Schille, Maike Sigg, Marc Weigelt, Andre Weise, Nicole Zubizarreta; Germany, Hanover Clinical Centre: Olga Kordonouri, Thomas Danne, Laura Galuschka, Carolin Kruse, Sarah Landsberg, Karin Lange, Erika Marquardt, Felix Reschke, Frank Roloff, Thekla von dem Berge, Jantje Weiskorn, Mareike Polier, Bianca Schmidt. Germany, Munich Clinical Centre: Anette G. Ziegler, Peter Achenbach, Melanie Bunk, Willi Grätz, Anna Hofelich, Elisabeth Huber, Melina Kaiser, Alexandra Käbl, Annika Kölln, Benjamin Marcus, Annette Munzinger, Jasmin Ohli, Claudia Ramminger, Franziska Reinmüller, Veronika Vollmuth, Christiane Winkler; Poland Clinical Centre: Agnieszka Szypowska, Mariusz Ohtarzewski, Sylwia Dybkowska, Lidia Groele, Dorota Owczarek, Katarzyna Popko, Adrianna Cieloch, Elżbieta Górka, Agnieszka Mroczek, Beata Zduńczyk, Anna Zych, Wiktoria Czerwińska, Natalia Dziedzic; Sweden Clinical Centre: Helena Elding Larsson, Markus Lundgren, Hanna Samuelsson, Sofie Alström Mortin, Rasmus Bennet, Charlotte Brundin, Susanne Dahlberg, Lina Fransson, Ida Jönsson, Hannah Nenonen, Anita Ramelius, Carina Törn, Ulrika Ulvenhag, Marielle Lindström, Kobra Rhamati, Malin Goldman Tsubarah, Falastin Salami; UK, Oxford Clinical Centre: John A Todd, Manu Vatish, Matthew Snape, Rachel E J Besser, Tabitha Wishlade, Eleni Fotaki, Fenella Roseman, Dorota Pietrzak, Jude Mossop, Sophia Hawkins, Yama F Mujajidi, Rachel White, Conor Whelan, Sophie Vernon, Owen Bendor-Samuel, Helen Ratcliffe, Alice Lelliott, Aparna Reddy, Patrick Bose, Edel Claire, Ian Smith, Nazia Taj; Oxford vaccine group; and JDRF/Wellcome Diabetes and Inflammation Laboratory Group.

**Contributors** AJ, VV, CW, A-GZ, ML, HEL, OK, TvdB, EEB, M-LZ, AH, MO, AS and KC were responsible for collection of the patients' data. AJ and AW performed the statistical analyses. AJ, MW and KC wrote the manuscript. VV, CW, A-GZ, ML, HEL, OK, TvdB, EEB, M-LZ, AH, MO, AS, RB and JAT revised the manuscript. KC is guarantor of this study.

**Funding** This work was supported by the POInT Study, supported by the Leona M and Harry B Helmsley Charitable Trust (Helmsley) (grants #2018PG-T1D023 (The Global Platform for Prevention of Autoimmune Diabetes (GPPAD)-03 Study: POInT-Primary Oral Insulin Trial) and #2018PG-T1D062 (Biobanking for the Primary Oral Insulin Trial)), by the Helmholtz Zentrum, Munich, Germany and the BMBF, Germany (grant#01KX1818). UK Clinical Centre is also supported by Wellcome (107212/Z/15/Z) and JDRF (5-SRA-2015-130-A-N). The German Clinical Centers Munich and Dresden are also supported by the Deutsches Zentrum für Diabetesforschung DZD. The authors A-GZ and EEB are inventors of a patent entitled 'Method for determining the risk to develop type 1 diabetes' (WO 2019/002364).

**Disclaimer** The funding organisations had no role in the design of the trial.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

**Patient consent for publication** Not required.

**Ethics approval** This study involves human participants and was approved by the local ethical committees and regulatory authorities of the Technische Universität München, Medical Faculty (326/17 Af), the Medical University of Warsaw (Instytucje Matki i Dziecka w Warszawie) (199/2017), the UK Health Research Authority (18/SC/0019), Onderzoek UZ/KU Leuven (S60711) and the Regionala etikprövningsnämnden i Lund (2017/918). Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. The data that support the findings of this study are available on request from the GPPAD- Coordinating Centre.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

An Jacobs <http://orcid.org/0000-0003-2556-4454>

Olga Kordonouri <http://orcid.org/0000-0001-9563-3537>

#### REFERENCES

- Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357:266–81.
- Martens PJ, Gysemans C, Verstuyf A, et al. Vitamin D's effect on immune function. *Nutrients* 2020;12:1248.
- Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab* 2011;96:1911–30.
- Ross AC, Manson JE, Abrams SA, et al. The 2011 dietary reference intakes for calcium and vitamin D: what dietetics practitioners need to know. *J Am Diet Assoc* 2011;111:524–7.
- Bacchetta J, Edouard T, Laverny G, et al. Vitamin D and calcium intakes in general pediatric populations: a French expert consensus paper. *Arch Pediatr* 2022;29:312–25.
- Bischoff-Ferrari HA, Shao A, Dawson-Hughes B, et al. Benefit-risk assessment of vitamin D supplementation. *Osteoporos Int* 2010;21:1121–32.
- Malabanan A, Veronikis IE, Holick MF. Redefining vitamin D insufficiency. *Lancet* 1998;351:805–6.
- Martens P-J, Centelles-Lodeiro J, Ellis D, et al. High serum vitamin D concentrations, induced via diet, trigger immune and intestinal microbiota alterations leading to type 1 diabetes protection in NOD mice. *Front Immunol* 2022;13:902678.
- Kumar J, Muntner P, Kaskel FJ, et al. Prevalence and associations of 25-Hydroxyvitamin D deficiency in US children: NHANES 2001–2004. *Pediatrics* 2009;124:e362–70.
- Holick MF. The vitamin D deficiency pandemic: approaches for diagnosis, treatment and prevention. *Rev Endocr Metab Disord* 2017;18:153–65.
- Veldman CM, Cantorna MT, DeLuca HF. Expression of 1,25-dihydroxyvitamin D(3) receptor in the immune system. *Arch Biochem Biophys* 2000;374:334–8.
- Prieti B, Treiber G, Pieber TR, et al. Vitamin D and immune function. *Nutrients* 2013;5:2502–21.
- Pozzilli P, Manfrini S, Crinò A, et al. Low levels of 25-Hydroxyvitamin D3 and 1,25-dihydroxyvitamin D3 in patients with newly diagnosed type 1 diabetes. *Horm Metab Res* 2005;37:680–3.
- Littorin B, Blom P, Schölin A, et al. Lower levels of plasma 25-Hydroxyvitamin D among young adults at diagnosis of autoimmune type 1 diabetes compared with control subjects: results from the nationwide diabetes incidence study in Sweden (DISS). *Diabetologia* 2006;49:2847–52.
- Manousaki D, Harroud A, Mitchell RE, et al. Vitamin D levels and risk of type 1 diabetes: a Mendelian randomization study. *PLoS Med* 2021;18:e1003624.
- Knip M, Åkerblom HK, Becker D, et al. Hydrolyzed infant formula and early beta-cell autoimmunity: a randomized clinical trial. *JAMA* 2014;311:2279–87.
- Baeke F, van Etten E, Gysemans C, et al. Vitamin D signaling in immune-mediated disorders: evolving insights and therapeutic opportunities. *Mol Aspects Med* 2008;29:376–87.
- Not Available. Vitamin D supplement in early childhood and risk for type I (insulin-dependent) diabetes mellitus. *Diabetologia* 1999;42:51–4.
- Silvis K, Aronsson CA, Liu X, et al. Maternal dietary supplement use and development of islet autoimmunity in the offspring: TEDDY study. *Pediatr Diabetes* 2019;20:86–92.
- Thorsen SU, Mårild K, Olsen SF, et al. Lack of association between maternal or neonatal vitamin D status and risk of childhood type 1 diabetes: a scandinavian case-cohort study. *Am J Epidemiol* 2018;187:1174–81.
- Carakushansky M, Patel P, Ben Khallouq BA, et al. Prevalence of vitamin D deficiency in children with type 1 diabetes mellitus. *Cureus* 2020;12:e7836.
- Ziegler A-G, Achenbach P, Berner R, et al. Oral insulin therapy for primary prevention of type 1 diabetes in infants with high genetic risk: the GPPAD-point (global platform for the prevention of autoimmune diabetes primary oral insulin trial) study protocol. *BMJ Open* 2019;9:e028578.
- Bonifacio E, Ziegler A-G, Klingensmith G, et al. Effects of high-dose oral insulin on immune responses in children at high risk for type 1 diabetes: the pre-POINT randomized clinical trial. *JAMA* 2015;313:1541.
- He LP, Song YX, Zhu T, et al. Progress in the relationship between vitamin D deficiency and the incidence of type 1 diabetes mellitus in children. *J Diabetes Res* 2022;2022:5953562.
- Nair R, Maseeh A. Vitamin D: the "sunshine" vitamin. *J Pharmacol Pharmacother* 2012;3:118–26.
- Miettinen ME, Niinistö S, Erlund I, et al. Serum 25-Hydroxyvitamin D concentration in childhood and risk of islet autoimmunity and type 1 diabetes: the TRIGR nested case-control ancillary study. *Diabetologia* 2020;63:780–7.
- Ozcan A, Kendirci M, Kondolot M, et al. Evaluation of vitamin D prophylaxis in 3-36-month-old infants and children. *J Pediatr Endocrinol Metab* 2017;30:543–9.
- Kasemsripitak S, Jaruratanasirikul S, Boonrusmee S, et al. Prevalence and risk factors for vitamin D insufficiency in 6-12-month-old infants: a cross-sectional study in Southern Thailand. *BMC Pediatr* 2022;22:729.
- Bravo P. Vitamin D deficiency and insufficiency in healthy infants receiving standard supplementation. *Andes Pediatr* 2022;93:773–4.
- Deruyter S, Van Biervliet S, De Guchteneere A. Response to vitamin D replacement therapy in obese children and adolescents with vitamin D deficiency: a randomized controlled trial. *J Pediatr Endocrinol Metab* 2023;36:458–65.
- Nwosu BU, Parajuli S, Jasmin G, et al. Ergocalciferol in new-onset type 1 diabetes: a randomized controlled trial. *J Endocr Soc* 2022;6:bvab179.